



Soils Map



State: **Iowa**
 County: **Warren**
 Location: **15-74N-22W**
 Township: **White Breast**
 Acres: **115.38**
 Date: **9/6/2016**



Area Symbol: IA181, Soil Area Version: 20

| Code | Soil Description | Acres | Percent of field | CSR2 Legend | Non-Irr Class *c | *i Corn | *i Soybeans | CSR2** | CS R |
|------------------|---|-------|------------------|-------------|------------------|---------|-------------|--------|------|
| 23C2 | Arispe silty clay loam, 5 to 9 percent slopes, moderately eroded | 27.17 | 23.5% | | IIIe | 156.8 | 45.5 | 62 | 62 |
| 131B | Pershing silt loam, 2 to 5 percent slopes | 20.52 | 17.8% | | IIIe | 195.2 | 56.6 | 70 | 72 |
| 822D2 | Lamoni silty clay loam, 9 to 14 percent slopes, moderately eroded | 16.01 | 13.9% | | IVe | 100.8 | 29.2 | 11 | 15 |
| 185D2 | Bauer silt loam, 9 to 14 percent slopes, moderately eroded | 13.82 | 12.0% | | VIe | 124.8 | 36.2 | 15 | 20 |
| 131C2 | Pershing silt loam, 5 to 9 percent slopes, moderately eroded | 12.19 | 10.6% | | IIIe | 169.6 | 49.2 | 62 | 50 |
| 980B | Gullied land-Ely-Colo complex, 2 to 5 percent slopes | 12.08 | 10.5% | | VIe | 88 | 25.5 | 42 | 25 |
| 94E2 | Mystic-Caleb complex, 14 to 18 percent slopes, moderately eroded | 9.05 | 7.8% | | VIe | 88 | 25.5 | 19 | 10 |
| 1313F | Munterville silt loam, 18 to 35 percent slopes | 2.55 | 2.2% | | VIe | 88 | 25.5 | 5 | 5 |
| 864B | Grundy silty clay loam, 2 to 5 percent slopes | 1.53 | 1.3% | | Ile | 198.4 | 57.5 | 72 | 82 |
| 369 | Winterset silty clay loam, 0 to 2 percent slopes | 0.46 | 0.4% | | IIw | 217.6 | 63.1 | 84 | 83 |
| Weighted Average | | | | | | 140.1 | 40.6 | 44.2 | 42.1 |

Area Symbol: IA181, Soil Area Version: 20

**IA has updated the CSR values for each county to CSR2.

*i Yield data provided by the ISPAID Database version 8.1 developed by IA State University.

*c: Using Capabilities Class Dominant Condition Aggregation Method

Soils data provided by USDA and NRCS.