



United States
Department of
Agriculture


NRCS

Natural
Resources
Conservation
Service

A product of the National
Cooperative Soil Survey,
a joint effort of the United
States Department of
Agriculture and other
Federal agencies, State
agencies including the
Agricultural Experiment
Stations, and local
participants

Custom Soil Resource Report for Benewah County Area, Idaho, Western Part; and Kootenai County Area, Idaho

Sunny Slopes Farm



Preface

Soil surveys contain information that affects land use planning in survey areas. They highlight soil limitations that affect various land uses and provide information about the properties of the soils in the survey areas. Soil surveys are designed for many different users, including farmers, ranchers, foresters, agronomists, urban planners, community officials, engineers, developers, builders, and home buyers. Also, conservationists, teachers, students, and specialists in recreation, waste disposal, and pollution control can use the surveys to help them understand, protect, or enhance the environment.

Various land use regulations of Federal, State, and local governments may impose special restrictions on land use or land treatment. Soil surveys identify soil properties that are used in making various land use or land treatment decisions. The information is intended to help the land users identify and reduce the effects of soil limitations on various land uses. The landowner or user is responsible for identifying and complying with existing laws and regulations.

Although soil survey information can be used for general farm, local, and wider area planning, onsite investigation is needed to supplement this information in some cases. Examples include soil quality assessments (<http://www.nrcs.usda.gov/wps/portal/nrcs/main/soils/health/>) and certain conservation and engineering applications. For more detailed information, contact your local USDA Service Center (<http://offices.sc.egov.usda.gov/locator/app?agency=nrcs>) or your NRCS State Soil Scientist (http://www.nrcs.usda.gov/wps/portal/nrcs/detail/soils/contactus/?cid=nrcs142p2_053951).

Great differences in soil properties can occur within short distances. Some soils are seasonally wet or subject to flooding. Some are too unstable to be used as a foundation for buildings or roads. Clayey or wet soils are poorly suited to use as septic tank absorption fields. A high water table makes a soil poorly suited to basements or underground installations.

The National Cooperative Soil Survey is a joint effort of the United States Department of Agriculture and other Federal agencies, State agencies including the Agricultural Experiment Stations, and local agencies. The Natural Resources Conservation Service (NRCS) has leadership for the Federal part of the National Cooperative Soil Survey.

Information about soils is updated periodically. Updated information is available through the NRCS Web Soil Survey, the site for official soil survey information.

The U.S. Department of Agriculture (USDA) prohibits discrimination in all its programs and activities on the basis of race, color, national origin, age, disability, and where applicable, sex, marital status, familial status, parental status, religion, sexual orientation, genetic information, political beliefs, reprisal, or because all or a part of an individual's income is derived from any public assistance program. (Not all prohibited bases apply to all programs.) Persons with disabilities who require alternative means

for communication of program information (Braille, large print, audiotape, etc.) should contact USDA's TARGET Center at (202) 720-2600 (voice and TDD). To file a complaint of discrimination, write to USDA, Director, Office of Civil Rights, 1400 Independence Avenue, S.W., Washington, D.C. 20250-9410 or call (800) 795-3272 (voice) or (202) 720-6382 (TDD). USDA is an equal opportunity provider and employer.

Contents

Preface	2
How Soil Surveys Are Made	5
Soil Map	7
Soil Map.....	8
Legend.....	9
Map Unit Legend.....	10
Map Unit Descriptions.....	11
Benewah County Area, Idaho, Western Part.....	13
121—Latahco-Lovell complex, 0 to 3 percent slopes.....	13
201—Blinn ashy silt loam, 35 to 65 percent slopes, stony.....	15
202—Blinn-Bobbitt complex, 35 to 65 percent slopes, stony.....	16
230—Lacy, stony-Rock outcrop complex, 5 to 35 percent slopes.....	17
231—Lacy, very stony-Rock outcrop complex, 35 to 65 percent slopes.....	19
232—Lacy-Bobbitt complex, 5 to 35 percent slopes, stony.....	20
233—Lacy-Bobbitt complex, 35 to 65 percent slopes, very stony.....	22
300—Taney ashy silt loam, 3 to 8 percent slopes.....	23
310—Santa ashy silt loam, 2 to 8 percent slopes.....	24
311—Santa ashy silt loam, 8 to 15 percent slopes.....	25
335—Carlinton ashy silt loam, dry, 8 to 25 percent slopes.....	26
336—Carlinton, dry-Taney complex, 3 to 8 percent slopes.....	28
Kootenai County Area, Idaho.....	30
1hn6j—Blinn ashy silt loam, 5 to 35 percent slopes, stony.....	30
135—Lacy gravelly loam, very stony-Rock outcrop complex, 5 to 35 percent slopes.....	31
146—McCrosket-Ardenvoir association, 20 to 35 percent slopes.....	32
157—Porrett silt loam.....	33
165—Santa ashy silt loam, 2 to 8 percent slopes.....	34
166—Santa ashy silt loam, 8 to 15 percent slopes.....	35
168—Santa variant silt loam, 5 to 20 percent slopes.....	36
186—Taney silt loam, 3 to 7 percent slopes.....	37
187—Taney silt loam, 7 to 25 percent slopes.....	38
189—Tekoa gravelly silt loam, 5 to 20 percent slopes.....	39
232—Lacy-Bobbitt complex, 5 to 35 percent slopes, stony.....	40
335—Carlinton ashy silt loam, dry, 8 to 25 percent slopes.....	42
336—Carlinton, dry-Taney complex, 3 to 8 percent slopes.....	43
References	45

How Soil Surveys Are Made

Soil surveys are made to provide information about the soils and miscellaneous areas in a specific area. They include a description of the soils and miscellaneous areas and their location on the landscape and tables that show soil properties and limitations affecting various uses. Soil scientists observed the steepness, length, and shape of the slopes; the general pattern of drainage; the kinds of crops and native plants; and the kinds of bedrock. They observed and described many soil profiles. A soil profile is the sequence of natural layers, or horizons, in a soil. The profile extends from the surface down into the unconsolidated material in which the soil formed or from the surface down to bedrock. The unconsolidated material is devoid of roots and other living organisms and has not been changed by other biological activity.

Currently, soils are mapped according to the boundaries of major land resource areas (MLRAs). MLRAs are geographically associated land resource units that share common characteristics related to physiography, geology, climate, water resources, soils, biological resources, and land uses (USDA, 2006). Soil survey areas typically consist of parts of one or more MLRA.

The soils and miscellaneous areas in a survey area occur in an orderly pattern that is related to the geology, landforms, relief, climate, and natural vegetation of the area. Each kind of soil and miscellaneous area is associated with a particular kind of landform or with a segment of the landform. By observing the soils and miscellaneous areas in the survey area and relating their position to specific segments of the landform, a soil scientist develops a concept, or model, of how they were formed. Thus, during mapping, this model enables the soil scientist to predict with a considerable degree of accuracy the kind of soil or miscellaneous area at a specific location on the landscape.

Commonly, individual soils on the landscape merge into one another as their characteristics gradually change. To construct an accurate soil map, however, soil scientists must determine the boundaries between the soils. They can observe only a limited number of soil profiles. Nevertheless, these observations, supplemented by an understanding of the soil-vegetation-landscape relationship, are sufficient to verify predictions of the kinds of soil in an area and to determine the boundaries.

Soil scientists recorded the characteristics of the soil profiles that they studied. They noted soil color, texture, size and shape of soil aggregates, kind and amount of rock fragments, distribution of plant roots, reaction, and other features that enable them to identify soils. After describing the soils in the survey area and determining their properties, the soil scientists assigned the soils to taxonomic classes (units). Taxonomic classes are concepts. Each taxonomic class has a set of soil characteristics with precisely defined limits. The classes are used as a basis for comparison to classify soils systematically. Soil taxonomy, the system of taxonomic classification used in the United States, is based mainly on the kind and character of soil properties and the arrangement of horizons within the profile. After the soil scientists classified and named the soils in the survey area, they compared the

individual soils with similar soils in the same taxonomic class in other areas so that they could confirm data and assemble additional data based on experience and research.

The objective of soil mapping is not to delineate pure map unit components; the objective is to separate the landscape into landforms or landform segments that have similar use and management requirements. Each map unit is defined by a unique combination of soil components and/or miscellaneous areas in predictable proportions. Some components may be highly contrasting to the other components of the map unit. The presence of minor components in a map unit in no way diminishes the usefulness or accuracy of the data. The delineation of such landforms and landform segments on the map provides sufficient information for the development of resource plans. If intensive use of small areas is planned, onsite investigation is needed to define and locate the soils and miscellaneous areas.

Soil scientists make many field observations in the process of producing a soil map. The frequency of observation is dependent upon several factors, including scale of mapping, intensity of mapping, design of map units, complexity of the landscape, and experience of the soil scientist. Observations are made to test and refine the soil-landscape model and predictions and to verify the classification of the soils at specific locations. Once the soil-landscape model is refined, a significantly smaller number of measurements of individual soil properties are made and recorded. These measurements may include field measurements, such as those for color, depth to bedrock, and texture, and laboratory measurements, such as those for content of sand, silt, clay, salt, and other components. Properties of each soil typically vary from one point to another across the landscape.

Observations for map unit components are aggregated to develop ranges of characteristics for the components. The aggregated values are presented. Direct measurements do not exist for every property presented for every map unit component. Values for some properties are estimated from combinations of other properties.

While a soil survey is in progress, samples of some of the soils in the area generally are collected for laboratory analyses and for engineering tests. Soil scientists interpret the data from these analyses and tests as well as the field-observed characteristics and the soil properties to determine the expected behavior of the soils under different uses. Interpretations for all of the soils are field tested through observation of the soils in different uses and under different levels of management. Some interpretations are modified to fit local conditions, and some new interpretations are developed to meet local needs. Data are assembled from other sources, such as research information, production records, and field experience of specialists. For example, data on crop yields under defined levels of management are assembled from farm records and from field or plot experiments on the same kinds of soil.


Predictions about soil behavior are based not only on soil properties but also on such variables as climate and biological activity. Soil conditions are predictable over long periods of time, but they are not predictable from year to year. For example, soil scientists can predict with a fairly high degree of accuracy that a given soil will have a high water table within certain depths in most years, but they cannot predict that a high water table will always be at a specific level in the soil on a specific date.

After soil scientists located and identified the significant natural bodies of soil in the survey area, they drew the boundaries of these bodies on aerial photographs and identified each as a specific map unit. Aerial photographs show trees, buildings, fields, roads, and rivers, all of which help in locating boundaries accurately.

Soil Map

The soil map section includes the soil map for the defined area of interest, a list of soil map units on the map and extent of each map unit, and cartographic symbols displayed on the map. Also presented are various metadata about data used to produce the map, and a description of each soil map unit.

Custom Soil Resource Report Soil Map



Custom Soil Resource Report

MAP LEGEND

Area of Interest (AOI)

 Area of Interest (AOI)

Soils

 Soil Map Unit Polygons


 Soil Map Unit Lines


 Soil Map Unit Points

Special Point Features

 Blowout

 Borrow Pit


 Clay Spot


 Closed Depression

 Gravel Pit

 Gravelly Spot


 Landfill

 Lava Flow

 Marsh or swamp

 Mine or Quarry


 Miscellaneous Water


 Perennial Water

 Rock Outcrop


 Saline Spot

 Sandy Spot

 Severely Eroded Spot


 Sinkhole


 Slide or Slip

 Sodic Spot


 Spoil Area

 Stony Spot

 Very Stony Spot

 Wet Spot

 Other

 Special Line Features

Water Features

 Streams and Canals


Transportation

 Rails

 Interstate Highways

 US Routes

 Major Roads

 Local Roads

Background

 Aerial Photography

MAP INFORMATION

The soil surveys that comprise your AOI were mapped at 1:24,000.

Please rely on the bar scale on each map sheet for map measurements.

Source of Map: Natural Resources Conservation Service
Web Soil Survey URL: <http://websoilsurvey.nrcs.usda.gov>
Coordinate System: Web Mercator (EPSG:3857)

Maps from the Web Soil Survey are based on the Web Mercator projection, which preserves direction and shape but distorts distance and area. A projection that preserves area, such as the Albers equal-area conic projection, should be used if more accurate calculations of distance or area are required.

This product is generated from the USDA-NRCS certified data as of the version date(s) listed below.

Soil Survey Area: Benewah County Area, Idaho, Western Part
Survey Area Data: Version 3, Sep 25, 2015

Soil Survey Area: Kootenai County Area, Idaho
Survey Area Data: Version 13, Sep 9, 2015

Your area of interest (AOI) includes more than one soil survey area. These survey areas may have been mapped at different scales, with a different land use in mind, at different times, or at different levels of detail. This may result in map unit symbols, soil properties, and interpretations that do not completely agree across soil survey area boundaries.

Soil map units are labeled (as space allows) for map scales 1:50,000 or larger.

Date(s) aerial images were photographed: Jul 5, 2011—Jul 29, 2011

The orthophoto or other base map on which the soil lines were compiled and digitized probably differs from the background imagery displayed on these maps. As a result, some minor shifting of map unit boundaries may be evident.

Map Unit Legend

Benewah County Area, Idaho, Western Part (ID620)			
Map Unit Symbol	Map Unit Name	Acres in AOI	Percent of AOI
121	Latahco-Lovell complex, 0 to 3 percent slopes	31.1	2.4%
201	Blinn ashy silt loam, 35 to 65 percent slopes, stony	12.0	0.9%
202	Blinn-Bobbitt complex, 35 to 65 percent slopes, stony	0.9	0.1%
230	Lacy, stony-Rock outcrop complex, 5 to 35 percent slopes	28.4	2.2%
231	Lacy, very stony-Rock outcrop complex, 35 to 65 percent slopes	21.6	1.7%
232	Lacy-Bobbitt complex, 5 to 35 percent slopes, stony	34.4	2.6%
233	Lacy-Bobbitt complex, 35 to 65 percent slopes, very stony	12.0	0.9%
300	Taney ashy silt loam, 3 to 8 percent slopes	16.2	1.2%
310	Santa ashy silt loam, 2 to 8 percent slopes	95.4	7.3%
311	Santa ashy silt loam, 8 to 15 percent slopes	25.2	1.9%
335	Carlinton ashy silt loam, dry, 8 to 25 percent slopes	219.4	16.8%
336	Carlinton, dry-Taney complex, 3 to 8 percent slopes	373.7	28.6%
Subtotals for Soil Survey Area		870.3	66.7%
Totals for Area of Interest		1,305.5	100.0%

Kootenai County Area, Idaho (ID606)			
Map Unit Symbol	Map Unit Name	Acres in AOI	Percent of AOI
1hn6j	Blinn ashy silt loam, 5 to 35 percent slopes, stony	7.6	0.6%
135	Lacy gravelly loam, very stony-Rock outcrop complex, 5 to 35 percent slopes	2.6	0.2%
146	McCrosket-Ardenvoir association, 20 to 35 percent slopes	0.7	0.1%
157	Porrett silt loam	32.1	2.5%
165	Santa ashy silt loam, 2 to 8 percent slopes	105.4	8.1%
166	Santa ashy silt loam, 8 to 15 percent slopes	84.8	6.5%

Custom Soil Resource Report

Kootenai County Area, Idaho (ID606)			
Map Unit Symbol	Map Unit Name	Acres in AOI	Percent of AOI
168	Santa variant silt loam, 5 to 20 percent slopes	13.8	1.1%
186	Taney silt loam, 3 to 7 percent slopes	9.4	0.7%
187	Taney silt loam, 7 to 25 percent slopes	47.6	3.6%
189	Tekoa gravelly silt loam, 5 to 20 percent slopes	38.3	2.9%
232	Lacy-Bobbitt complex, 5 to 35 percent slopes, stony	4.1	0.3%
335	Carlinton ashy silt loam, dry, 8 to 25 percent slopes	30.1	2.3%
336	Carlinton, dry-Taney complex, 3 to 8 percent slopes	58.8	4.5%
Subtotals for Soil Survey Area		435.2	33.3%
Totals for Area of Interest		1,305.5	100.0%

Map Unit Descriptions

The map units delineated on the detailed soil maps in a soil survey represent the soils or miscellaneous areas in the survey area. The map unit descriptions, along with the maps, can be used to determine the composition and properties of a unit.

A map unit delineation on a soil map represents an area dominated by one or more major kinds of soil or miscellaneous areas. A map unit is identified and named according to the taxonomic classification of the dominant soils. Within a taxonomic class there are precisely defined limits for the properties of the soils. On the landscape, however, the soils are natural phenomena, and they have the characteristic variability of all natural phenomena. Thus, the range of some observed properties may extend beyond the limits defined for a taxonomic class. Areas of soils of a single taxonomic class rarely, if ever, can be mapped without including areas of other taxonomic classes. Consequently, every map unit is made up of the soils or miscellaneous areas for which it is named and some minor components that belong to taxonomic classes other than those of the major soils.

Most minor soils have properties similar to those of the dominant soil or soils in the map unit, and thus they do not affect use and management. These are called noncontrasting, or similar, components. They may or may not be mentioned in a particular map unit description. Other minor components, however, have properties and behavioral characteristics divergent enough to affect use or to require different management. These are called contrasting, or dissimilar, components. They generally are in small areas and could not be mapped separately because of the scale used. Some small areas of strongly contrasting soils or miscellaneous areas are identified by a special symbol on the maps. If included in the database for a given area, the contrasting minor components are identified in the map unit descriptions along with some characteristics of each. A few areas of minor components may not have been observed, and consequently they are not mentioned in the descriptions, especially

where the pattern was so complex that it was impractical to make enough observations to identify all the soils and miscellaneous areas on the landscape.

The presence of minor components in a map unit in no way diminishes the usefulness or accuracy of the data. The objective of mapping is not to delineate pure taxonomic classes but rather to separate the landscape into landforms or landform segments that have similar use and management requirements. The delineation of such segments on the map provides sufficient information for the development of resource plans. If intensive use of small areas is planned, however, onsite investigation is needed to define and locate the soils and miscellaneous areas.

An identifying symbol precedes the map unit name in the map unit descriptions. Each description includes general facts about the unit and gives important soil properties and qualities.

Soils that have profiles that are almost alike make up a *soil series*. Except for differences in texture of the surface layer, all the soils of a series have major horizons that are similar in composition, thickness, and arrangement.

Soils of one series can differ in texture of the surface layer, slope, stoniness, salinity, degree of erosion, and other characteristics that affect their use. On the basis of such differences, a soil series is divided into *soil phases*. Most of the areas shown on the detailed soil maps are phases of soil series. The name of a soil phase commonly indicates a feature that affects use or management. For example, Alpha silt loam, 0 to 2 percent slopes, is a phase of the Alpha series.

Some map units are made up of two or more major soils or miscellaneous areas. These map units are complexes, associations, or undifferentiated groups.

A *complex* consists of two or more soils or miscellaneous areas in such an intricate pattern or in such small areas that they cannot be shown separately on the maps. The pattern and proportion of the soils or miscellaneous areas are somewhat similar in all areas. Alpha-Beta complex, 0 to 6 percent slopes, is an example.

An *association* is made up of two or more geographically associated soils or miscellaneous areas that are shown as one unit on the maps. Because of present or anticipated uses of the map units in the survey area, it was not considered practical or necessary to map the soils or miscellaneous areas separately. The pattern and relative proportion of the soils or miscellaneous areas are somewhat similar. Alpha-Beta association, 0 to 2 percent slopes, is an example.

An *undifferentiated group* is made up of two or more soils or miscellaneous areas that could be mapped individually but are mapped as one unit because similar interpretations can be made for use and management. The pattern and proportion of the soils or miscellaneous areas in a mapped area are not uniform. An area can be made up of only one of the major soils or miscellaneous areas, or it can be made up of all of them. Alpha and Beta soils, 0 to 2 percent slopes, is an example.

Some surveys include *miscellaneous areas*. Such areas have little or no soil material and support little or no vegetation. Rock outcrop is an example.

Benewah County Area, Idaho, Western Part

121—Latahco-Lovell complex, 0 to 3 percent slopes

Map Unit Setting

National map unit symbol: 1hn5c
Elevation: 2,500 to 2,890 feet
Mean annual precipitation: 22 to 28 inches
Mean annual air temperature: 41 to 46 degrees F
Frost-free period: 90 to 120 days
Farmland classification: Prime farmland if drained

Map Unit Composition

Latahco and similar soils: 60 percent
Lovell and similar soils: 30 percent
Minor components: 8 percent
Estimates are based on observations, descriptions, and transects of the mapunit.

Description of Latahco

Setting

Landform: Terraces, drainageways
Landform position (two-dimensional): Toeslope
Down-slope shape: Linear
Across-slope shape: Convex
Parent material: Loess

Typical profile

Ap - 0 to 13 inches: silt loam
E - 13 to 20 inches: silt loam
Bt - 20 to 26 inches: silty clay loam
Btk - 26 to 42 inches: silty clay loam
Btb - 42 to 51 inches: silt loam
Cc - 51 to 62 inches: silt loam

Properties and qualities

Slope: 0 to 3 percent
Depth to restrictive feature: More than 80 inches
Natural drainage class: Somewhat poorly drained
Capacity of the most limiting layer to transmit water (Ksat): Moderately high to high
(0.57 to 1.98 in/hr)
Depth to water table: About 16 to 21 inches
Frequency of flooding: Occasional
Frequency of ponding: None
Calcium carbonate, maximum in profile: 4 percent
Available water storage in profile: High (about 11.0 inches)

Interpretive groups

Land capability classification (irrigated): 3w
Land capability classification (nonirrigated): 3w
Hydrologic Soil Group: B/D
Ecological site: DRY MEADOW (R009XY019ID)

Description of Lovell

Setting

Landform: Flood plains
Landform position (two-dimensional): Toeslope
Landform position (three-dimensional): Riser
Down-slope shape: Concave
Across-slope shape: Linear
Parent material: Volcanic ash and loess over alluvium

Typical profile

Ap - 0 to 8 inches: ashy silt loam
Eg - 8 to 18 inches: ashy silt loam
EBtg - 18 to 22 inches: silt loam
Bt1 - 22 to 34 inches: silt loam
2Bt2 - 34 to 51 inches: loam
2BC - 51 to 60 inches: loam

Properties and qualities

Slope: 0 to 2 percent
Depth to restrictive feature: More than 80 inches
Natural drainage class: Poorly drained
Capacity of the most limiting layer to transmit water (Ksat): Moderately high (0.20 to 0.57 in/hr)
Depth to water table: About 8 to 26 inches
Frequency of flooding: Occasional
Frequency of ponding: None
Available water storage in profile: High (about 10.6 inches)

Interpretive groups

Land capability classification (irrigated): 5w
Land capability classification (nonirrigated): 5w
Hydrologic Soil Group: C/D
Ecological site: MEADOW (R009XY018ID)

Minor Components

Endoaquolls

Percent of map unit: 8 percent
Landform: Flood plains, drainageways
Landform position (two-dimensional): Toeslope
Landform position (three-dimensional): Tread
Down-slope shape: Concave
Across-slope shape: Concave
Ecological site: WET MEADOW 16-24 PZ (R009XY601WA)

201—Blinn ashy silt loam, 35 to 65 percent slopes, stony

Map Unit Setting

National map unit symbol: 1hn6k
Elevation: 2,100 to 3,000 feet
Mean annual precipitation: 25 to 30 inches
Mean annual air temperature: 43 to 45 degrees F
Frost-free period: 90 to 110 days
Farmland classification: Not prime farmland

Map Unit Composition

Blinn, stony surface, and similar soils: 80 percent
Estimates are based on observations, descriptions, and transects of the mapunit.

Description of Blinn, Stony Surface

Setting

Landform: Escarpments, canyons
Landform position (two-dimensional): Backslope
Landform position (three-dimensional): Side slope
Down-slope shape: Convex
Across-slope shape: Convex
Parent material: Volcanic ash and loess over colluvium over residuum weathered from basalt

Typical profile

Oi - 0 to 1 inches: slightly decomposed plant material
Oe - 1 to 2 inches: moderately decomposed plant material
A - 2 to 6 inches: ashy silt loam
Bw1 - 6 to 12 inches: gravelly ashy silt loam
Bw2 - 12 to 24 inches: stony loam
C - 24 to 39 inches: very stony loam
R - 39 to 40 inches: bedrock

Properties and qualities

Slope: 35 to 65 percent
Percent of area covered with surface fragments: 0.1 percent
Depth to restrictive feature: 20 to 40 inches to lithic bedrock
Natural drainage class: Well drained
Capacity of the most limiting layer to transmit water (Ksat): Moderately high to high (0.57 to 1.98 in/hr)
Depth to water table: More than 80 inches
Frequency of flooding: None
Frequency of ponding: None
Available water storage in profile: Low (about 5.1 inches)

Interpretive groups

Land capability classification (irrigated): None specified
Land capability classification (nonirrigated): 7e

Custom Soil Resource Report

Hydrologic Soil Group: C

Other vegetative classification: grand fir/ninebark (CN506)

202—Blinn-Bobbitt complex, 35 to 65 percent slopes, stony

Map Unit Setting

National map unit symbol: 1hn6n

Elevation: 2,100 to 3,000 feet

Mean annual precipitation: 25 to 30 inches

Mean annual air temperature: 43 to 49 degrees F

Frost-free period: 95 to 130 days

Farmland classification: Not prime farmland

Map Unit Composition

Blinn, stony surface, and similar soils: 55 percent

Bobbitt, stony surface, and similar soils: 30 percent

Estimates are based on observations, descriptions, and transects of the mapunit.

Description of Blinn, Stony Surface

Setting

Landform: Escarpments, canyons

Landform position (two-dimensional): Backslope

Landform position (three-dimensional): Side slope

Down-slope shape: Convex

Across-slope shape: Convex

Parent material: Volcanic ash and loess over colluvium over residuum weathered from basalt

Typical profile

O_i - 0 to 1 inches: slightly decomposed plant material

O_e - 1 to 2 inches: moderately decomposed plant material

A - 2 to 6 inches: ashy silt loam

Bw₁ - 6 to 12 inches: gravelly ashy silt loam

Bw₂ - 12 to 24 inches: stony loam

C - 24 to 39 inches: very stony loam

R - 39 to 40 inches: bedrock

Properties and qualities

Slope: 35 to 65 percent

Percent of area covered with surface fragments: 0.1 percent

Depth to restrictive feature: 20 to 40 inches to lithic bedrock

Natural drainage class: Well drained

Capacity of the most limiting layer to transmit water (K_{sat}): Moderately high to high (0.57 to 1.98 in/hr)

Depth to water table: More than 80 inches

Frequency of flooding: None

Frequency of ponding: None

Available water storage in profile: Low (about 5.1 inches)

Interpretive groups

Land capability classification (irrigated): None specified

Custom Soil Resource Report

Land capability classification (nonirrigated): 7e

Hydrologic Soil Group: C

Other vegetative classification: grand fir/ninebark (CN506)

Description of Bobbitt, Stony Surface

Setting

Landform: Escarpments, canyons

Landform position (two-dimensional): Backslope

Landform position (three-dimensional): Side slope

Down-slope shape: Convex

Across-slope shape: Convex

Parent material: Volcanic ash and loess over colluvium over residuum weathered from basalt

Typical profile

Oi - 0 to 1 inches: slightly decomposed plant material

Oe - 1 to 2 inches: moderately decomposed plant material

A - 2 to 9 inches: stony ashy silt loam

Bt - 9 to 23 inches: very stony clay loam

R - 23 to 33 inches: bedrock

Properties and qualities

Slope: 35 to 65 percent

Percent of area covered with surface fragments: 0.1 percent

Depth to restrictive feature: 20 to 40 inches to lithic bedrock

Natural drainage class: Well drained

Capacity of the most limiting layer to transmit water (Ksat): Moderately high to high (0.57 to 1.98 in/hr)

Depth to water table: More than 80 inches

Frequency of flooding: None

Frequency of ponding: None

Available water storage in profile: Low (about 3.3 inches)

Interpretive groups

Land capability classification (irrigated): None specified

Land capability classification (nonirrigated): 7e

Hydrologic Soil Group: C

Other vegetative classification: Douglas-fir/ninebark (CN260)

230—Lacy, stony-Rock outcrop complex, 5 to 35 percent slopes

Map Unit Setting

National map unit symbol: 1hn6q

Elevation: 2,130 to 2,950 feet

Mean annual precipitation: 25 to 30 inches

Mean annual air temperature: 47 to 50 degrees F

Frost-free period: 100 to 140 days

Farmland classification: Not prime farmland

Map Unit Composition

Lacy, stony surface, and similar soils: 65 percent

Rock outcrop: 15 percent

Estimates are based on observations, descriptions, and transects of the mapunit.

Description of Lacy, Stony Surface

Setting

Landform: Escarpments, structural benches

Landform position (two-dimensional): Backslope, summit, shoulder

Landform position (three-dimensional): Side slope

Down-slope shape: Convex

Across-slope shape: Convex

Parent material: Loess over colluvium over bedrock derived from basalt

Typical profile

Oi - 0 to 1 inches: slightly decomposed plant material

Oe - 1 to 2 inches: moderately decomposed plant material

A1 - 2 to 3 inches: stony silt loam

A2 - 3 to 10 inches: stony silt loam

Bt1 - 10 to 14 inches: very stony silt loam

Bt2 - 14 to 17 inches: extremely stony loam

R - 17 to 27 inches: bedrock

Properties and qualities

Slope: 5 to 35 percent

Percent of area covered with surface fragments: 0.1 percent

Depth to restrictive feature: 10 to 20 inches to lithic bedrock

Natural drainage class: Well drained

Capacity of the most limiting layer to transmit water (Ksat): Moderately high to high
(0.57 to 1.98 in/hr)

Depth to water table: More than 80 inches

Frequency of flooding: None

Frequency of ponding: None

Available water storage in profile: Very low (about 2.5 inches)

Interpretive groups

Land capability classification (irrigated): None specified

Land capability classification (nonirrigated): 6e

Hydrologic Soil Group: D

Other vegetative classification: ponderosa pine/common snowberry (CN170)

Description of Rock Outcrop

Typical profile

R - 0 to 60 inches: bedrock

Properties and qualities

Slope: 5 to 35 percent

Depth to restrictive feature: 0 inches to lithic bedrock

Interpretive groups

Land capability classification (irrigated): None specified

Land capability classification (nonirrigated): 8

231—Lacy, very stony-Rock outcrop complex, 35 to 65 percent slopes

Map Unit Setting

National map unit symbol: 1hn6r
Elevation: 2,150 to 3,000 feet
Mean annual precipitation: 25 to 30 inches
Mean annual air temperature: 47 to 50 degrees F
Frost-free period: 100 to 140 days
Farmland classification: Not prime farmland

Map Unit Composition

Lacy, very stony surface, and similar soils: 60 percent
Rock outcrop: 25 percent
Estimates are based on observations, descriptions, and transects of the mapunit.

Description of Lacy, Very Stony Surface

Setting

Landform: Escarpments, canyons
Landform position (two-dimensional): Backslope
Landform position (three-dimensional): Side slope
Down-slope shape: Convex
Across-slope shape: Convex
Parent material: Loess over colluvium over bedrock derived from basalt

Typical profile

Oi - 0 to 1 inches: slightly decomposed plant material
A1 - 1 to 2 inches: stony loam
A2 - 2 to 4 inches: stony loam
AB - 4 to 8 inches: very stony loam
Bt1 - 8 to 16 inches: very stony loam
Bt2 - 16 to 19 inches: extremely stony clay loam
R - 19 to 29 inches: bedrock

Properties and qualities

Slope: 35 to 65 percent
Percent of area covered with surface fragments: 1.0 percent
Depth to restrictive feature: 10 to 20 inches to lithic bedrock
Natural drainage class: Well drained
Capacity of the most limiting layer to transmit water (Ksat): Moderately high to high
(0.57 to 1.98 in/hr)
Depth to water table: More than 80 inches
Frequency of flooding: None
Frequency of ponding: None
Available water storage in profile: Very low (about 1.7 inches)

Interpretive groups

Land capability classification (irrigated): None specified
Land capability classification (nonirrigated): 7e
Hydrologic Soil Group: D

Other vegetative classification: ponderosa pine/common snowberry (CN170)

Description of Rock Outcrop

Typical profile

R - 0 to 60 inches: bedrock

Properties and qualities

Slope: 35 to 65 percent

Depth to restrictive feature: 0 inches to lithic bedrock

Interpretive groups

Land capability classification (irrigated): None specified

Land capability classification (nonirrigated): 8

232—Lacy-Bobbitt complex, 5 to 35 percent slopes, stony

Map Unit Setting

National map unit symbol: 1hn6s

Elevation: 2,130 to 3,000 feet

Mean annual precipitation: 25 to 30 inches

Mean annual air temperature: 47 to 50 degrees F

Frost-free period: 100 to 130 days

Farmland classification: Not prime farmland

Map Unit Composition

Lacy, stony surface, and similar soils: 55 percent

Bobbitt, stony surface, and similar soils: 30 percent

Estimates are based on observations, descriptions, and transects of the mapunit.

Description of Lacy, Stony Surface

Setting

Landform: Escarpments, structural benches

Landform position (two-dimensional): Backslope, shoulder

Landform position (three-dimensional): Side slope

Down-slope shape: Convex

Across-slope shape: Convex

Parent material: Loess over colluvium over bedrock derived from basalt

Typical profile

Oi - 0 to 1 inches: slightly decomposed plant material

Oe - 1 to 2 inches: moderately decomposed plant material

A1 - 2 to 3 inches: stony silt loam

A2 - 3 to 10 inches: stony silt loam

Bt1 - 10 to 14 inches: very stony silt loam

Bt2 - 14 to 17 inches: extremely stony loam

R - 17 to 27 inches: bedrock

Properties and qualities

Slope: 5 to 35 percent

Percent of area covered with surface fragments: 0.1 percent

Custom Soil Resource Report

Depth to restrictive feature: 10 to 20 inches to lithic bedrock

Natural drainage class: Well drained

Capacity of the most limiting layer to transmit water (Ksat): Moderately high to high
(0.57 to 1.98 in/hr)

Depth to water table: More than 80 inches

Frequency of flooding: None

Frequency of ponding: None

Available water storage in profile: Very low (about 2.5 inches)

Interpretive groups

Land capability classification (irrigated): None specified

Land capability classification (nonirrigated): 6e

Hydrologic Soil Group: D

Other vegetative classification: ponderosa pine/common snowberry (CN170)

Description of Bobbitt, Stony Surface

Setting

Landform: Escarpments, structural benches

Landform position (two-dimensional): Backslope, shoulder

Landform position (three-dimensional): Side slope

Down-slope shape: Convex

Across-slope shape: Convex

Parent material: Volcanic ash and loess over colluvium over residuum weathered from basalt

Typical profile

Oi - 0 to 1 inches: slightly decomposed plant material

Oe - 1 to 2 inches: moderately decomposed plant material

A - 2 to 9 inches: stony ashy silt loam

Bt - 9 to 23 inches: very stony clay loam

R - 23 to 33 inches: bedrock

Properties and qualities

Slope: 5 to 35 percent

Percent of area covered with surface fragments: 0.1 percent

Depth to restrictive feature: 20 to 40 inches to lithic bedrock

Natural drainage class: Well drained

Capacity of the most limiting layer to transmit water (Ksat): Moderately high to high
(0.57 to 1.98 in/hr)

Depth to water table: More than 80 inches

Frequency of flooding: None

Frequency of ponding: None

Available water storage in profile: Low (about 3.3 inches)

Interpretive groups

Land capability classification (irrigated): None specified

Land capability classification (nonirrigated): 4e

Hydrologic Soil Group: C

Other vegetative classification: Douglas-fir/ninebark (CN260)

233—Lacy-Bobbitt complex, 35 to 65 percent slopes, very stony

Map Unit Setting

National map unit symbol: 1hn6t
Elevation: 2,120 to 3,100 feet
Mean annual precipitation: 25 to 30 inches
Mean annual air temperature: 47 to 50 degrees F
Frost-free period: 100 to 130 days
Farmland classification: Not prime farmland

Map Unit Composition

Lacy, very stony surface, and similar soils: 55 percent
Bobbitt, very stony surface, and similar soils: 30 percent
Estimates are based on observations, descriptions, and transects of the mapunit.

Description of Lacy, Very Stony Surface

Setting

Landform: Escarpments, canyons
Landform position (two-dimensional): Backslope
Landform position (three-dimensional): Side slope
Down-slope shape: Convex
Across-slope shape: Convex
Parent material: Loess over colluvium over bedrock derived from basalt

Typical profile

Oi - 0 to 1 inches: slightly decomposed plant material
A1 - 1 to 2 inches: stony loam
A2 - 2 to 4 inches: stony loam
AB - 4 to 8 inches: very stony loam
Bt1 - 8 to 16 inches: very stony loam
Bt2 - 16 to 19 inches: extremely stony clay loam
R - 19 to 29 inches: bedrock

Properties and qualities

Slope: 35 to 65 percent
Percent of area covered with surface fragments: 1.0 percent
Depth to restrictive feature: 10 to 20 inches to lithic bedrock
Natural drainage class: Well drained
Capacity of the most limiting layer to transmit water (Ksat): Moderately high to high
(0.57 to 1.98 in/hr)
Depth to water table: More than 80 inches
Frequency of flooding: None
Frequency of ponding: None
Available water storage in profile: Very low (about 1.7 inches)

Interpretive groups

Land capability classification (irrigated): None specified
Land capability classification (nonirrigated): 7e
Hydrologic Soil Group: D

Other vegetative classification: ponderosa pine/common snowberry (CN170)

Description of Bobbitt, Very Stony Surface

Setting

Landform: Escarpments, canyons

Landform position (two-dimensional): Backslope

Landform position (three-dimensional): Side slope

Down-slope shape: Convex

Across-slope shape: Convex

Parent material: Volcanic ash and loess over colluvium over residuum weathered from basalt

Typical profile

Oi - 0 to 1 inches: slightly decomposed plant material

Oe - 1 to 2 inches: moderately decomposed plant material

A - 2 to 4 inches: stony ashy loam

AB - 4 to 11 inches: stony ashy loam

Bt1 - 11 to 15 inches: very cobbly loam

Bt2 - 15 to 27 inches: very cobbly loam

Bt3 - 27 to 33 inches: extremely stony loam

R - 33 to 43 inches: bedrock

Properties and qualities

Slope: 35 to 65 percent

Percent of area covered with surface fragments: 1.0 percent

Depth to restrictive feature: 20 to 40 inches to lithic bedrock

Natural drainage class: Well drained

Capacity of the most limiting layer to transmit water (Ksat): Moderately high to high (0.57 to 1.98 in/hr)

Depth to water table: More than 80 inches

Frequency of flooding: None

Frequency of ponding: None

Available water storage in profile: Low (about 4.1 inches)

Interpretive groups

Land capability classification (irrigated): None specified

Land capability classification (nonirrigated): 7e

Hydrologic Soil Group: C

Other vegetative classification: Douglas-fir/ninebark (CN260)

300—Taney ashy silt loam, 3 to 8 percent slopes

Map Unit Setting

National map unit symbol: 1hn70

Elevation: 2,560 to 3,210 feet

Mean annual precipitation: 25 to 28 inches

Mean annual air temperature: 43 to 45 degrees F

Frost-free period: 90 to 110 days

Farmland classification: Prime farmland if drained

Map Unit Composition

Taney and similar soils: 80 percent

Estimates are based on observations, descriptions, and transects of the mapunit.

Description of Taney

Setting

Landform: Loess hills

Landform position (two-dimensional): Shoulder, summit

Landform position (three-dimensional): Side slope

Down-slope shape: Concave

Across-slope shape: Linear

Parent material: Volcanic ash over loess

Typical profile

Oi - 0 to 1 inches: slightly decomposed plant material

Oe - 1 to 2 inches: moderately decomposed plant material

A - 2 to 4 inches: ashy silt loam

BA - 4 to 15 inches: ashy silt loam

Bw - 15 to 22 inches: silt loam

Bt - 22 to 29 inches: silt loam

EBc - 29 to 31 inches: silt loam

Btxcb - 31 to 53 inches: silty clay loam

Btxb - 53 to 60 inches: silty clay loam

Properties and qualities

Slope: 3 to 8 percent

Depth to restrictive feature: 23 to 40 inches to fragipan

Natural drainage class: Moderately well drained

Capacity of the most limiting layer to transmit water (Ksat): Moderately high to high
(0.57 to 1.98 in/hr)

Depth to water table: About 16 to 22 inches

Frequency of flooding: None

Frequency of ponding: None

Available water storage in profile: Moderate (about 6.7 inches)

Interpretive groups

Land capability classification (irrigated): 3e

Land capability classification (nonirrigated): 3e

Hydrologic Soil Group: C/D

Other vegetative classification: Douglas-fir/ninebark (CN260)

310—Santa ashy silt loam, 2 to 8 percent slopes

Map Unit Setting

National map unit symbol: 2v71x

Elevation: 2,610 to 3,200 feet

Mean annual precipitation: 25 to 35 inches

Mean annual air temperature: 43 to 46 degrees F

Frost-free period: 110 to 120 days

Farmland classification: Farmland of statewide importance, if drained

Map Unit Composition

Santa and similar soils: 85 percent

Estimates are based on observations, descriptions, and transects of the mapunit.

Description of Santa

Setting

Landform: Interfluves, hillslopes

Landform position (two-dimensional): Summit, shoulder, backslope

Landform position (three-dimensional): Interfluve, side slope

Down-slope shape: Convex, concave

Across-slope shape: Linear, convex

Parent material: Mixed volcanic ash and loess

Typical profile

Ap - 0 to 8 inches: ashy silt loam

Bw - 8 to 19 inches: silt loam

E - 19 to 29 inches: silt loam

Btxb1 - 29 to 38 inches: silt loam

Btxb2 - 38 to 59 inches: silt loam

Properties and qualities

Slope: 2 to 8 percent

Depth to restrictive feature: 24 to 39 inches to fragipan

Natural drainage class: Moderately well drained

Capacity of the most limiting layer to transmit water (Ksat): Very low to moderately low (0.00 to 0.06 in/hr)

Depth to water table: About 8 to 22 inches

Frequency of flooding: None

Frequency of ponding: None

Available water storage in profile: Low (about 5.7 inches)

Interpretive groups

Land capability classification (irrigated): None specified

Land capability classification (nonirrigated): 4e

Hydrologic Soil Group: C/D

Other vegetative classification: grand fir/ninebark (CN506)

311—Santa ashy silt loam, 8 to 15 percent slopes

Map Unit Setting

National map unit symbol: 2v71y

Elevation: 2,510 to 3,200 feet

Mean annual precipitation: 25 to 37 inches

Mean annual air temperature: 43 to 46 degrees F

Frost-free period: 105 to 120 days

Farmland classification: Farmland of statewide importance, if drained

Map Unit Composition

Santa and similar soils: 80 percent

Estimates are based on observations, descriptions, and transects of the mapunit.

Description of Santa

Setting

Landform: Interfluves, hillslopes

Landform position (two-dimensional): Summit, shoulder, backslope

Landform position (three-dimensional): Interfluve, side slope

Down-slope shape: Convex, concave

Across-slope shape: Linear, convex

Parent material: Mixed volcanic ash and loess

Typical profile

Ap - 0 to 8 inches: ashy silt loam

Bw - 8 to 19 inches: silt loam

E - 19 to 29 inches: silt loam

Btxb1 - 29 to 38 inches: silt loam

Btxb2 - 38 to 59 inches: silt loam

Properties and qualities

Slope: 8 to 15 percent

Depth to restrictive feature: 24 to 39 inches to fragipan

Natural drainage class: Moderately well drained

Capacity of the most limiting layer to transmit water (Ksat): Very low to moderately low (0.00 to 0.06 in/hr)

Depth to water table: About 8 to 22 inches

Frequency of flooding: None

Frequency of ponding: None

Available water storage in profile: Low (about 5.7 inches)

Interpretive groups

Land capability classification (irrigated): None specified

Land capability classification (nonirrigated): 4e

Hydrologic Soil Group: C/D

Other vegetative classification: grand fir/ninebark (CN506)

335—Carlinton ashy silt loam, dry, 8 to 25 percent slopes

Map Unit Setting

National map unit symbol: 1hn7g

Elevation: 2,700 to 3,230 feet

Mean annual precipitation: 25 to 28 inches

Mean annual air temperature: 43 to 45 degrees F

Frost-free period: 90 to 120 days

Farmland classification: Farmland of statewide importance, if drained

Map Unit Composition

Carlinton, dry, and similar soils: 80 percent

Minor components: 2 percent

Estimates are based on observations, descriptions, and transects of the mapunit.

Description of Carlinton, Dry

Setting

Landform: Loess hills
Landform position (two-dimensional): Backslope
Landform position (three-dimensional): Side slope
Down-slope shape: Convex
Across-slope shape: Linear
Parent material: Volcanic ash over loess

Typical profile

Ap1 - 0 to 5 inches: ashy silt loam
Ap2 - 5 to 10 inches: ashy silt loam
Bw - 10 to 14 inches: silt loam
EBt - 14 to 20 inches: silt loam
E - 20 to 23 inches: silt loam
BtbxE - 23 to 30 inches: silt loam
Btbx - 30 to 53 inches: silty clay loam
Btb - 53 to 60 inches: silty clay loam

Properties and qualities

Slope: 8 to 25 percent
Depth to restrictive feature: 21 to 40 inches to fragipan
Natural drainage class: Moderately well drained
Capacity of the most limiting layer to transmit water (Ksat): Moderately high to high (0.57 to 1.98 in/hr)
Depth to water table: About 14 to 20 inches
Frequency of flooding: None
Frequency of ponding: None
Available water storage in profile: Low (about 5.7 inches)

Interpretive groups

Land capability classification (irrigated): None specified
Land capability classification (nonirrigated): 3e
Hydrologic Soil Group: C/D
Other vegetative classification: Douglas-fir/ninebark (CN260)

Minor Components

Lovell

Percent of map unit: 2 percent
Landform: Drainageways
Landform position (two-dimensional): Toeslope
Down-slope shape: Concave
Across-slope shape: Linear
Ecological site: MEADOW (R009XY018ID)

336—Carlinton, dry-Taney complex, 3 to 8 percent slopes

Map Unit Setting

National map unit symbol: 1hn7h
Elevation: 2,560 to 3,020 feet
Mean annual precipitation: 25 to 28 inches
Mean annual air temperature: 43 to 45 degrees F
Frost-free period: 90 to 120 days
Farmland classification: Prime farmland if drained

Map Unit Composition

Carlinton, dry, and similar soils: 55 percent
Taney and similar soils: 25 percent
Estimates are based on observations, descriptions, and transects of the mapunit.

Description of Carlinton, Dry

Setting

Landform: Loess hills
Landform position (two-dimensional): Shoulder
Landform position (three-dimensional): Side slope
Down-slope shape: Convex
Across-slope shape: Linear
Parent material: Volcanic ash over loess

Typical profile

Ap1 - 0 to 5 inches: ashy silt loam
Ap2 - 5 to 10 inches: ashy silt loam
Bw - 10 to 14 inches: silt loam
EBt - 14 to 20 inches: silt loam
E - 20 to 23 inches: silt loam
BtbxE - 23 to 30 inches: silt loam
Btbx - 30 to 53 inches: silty clay loam
Btb - 53 to 60 inches: silty clay loam

Properties and qualities

Slope: 3 to 8 percent
Depth to restrictive feature: 21 to 40 inches to fragipan
Natural drainage class: Moderately well drained
Capacity of the most limiting layer to transmit water (Ksat): Moderately high to high
(0.57 to 1.98 in/hr)
Depth to water table: About 14 to 20 inches
Frequency of flooding: None
Frequency of ponding: None
Available water storage in profile: Low (about 5.7 inches)

Interpretive groups

Land capability classification (irrigated): 3e
Land capability classification (nonirrigated): 3e
Hydrologic Soil Group: C/D

Custom Soil Resource Report

Other vegetative classification: Douglas-fir/ninebark (CN260)

Description of Taney

Setting

Landform: Loess hills

Landform position (two-dimensional): Shoulder, summit

Landform position (three-dimensional): Side slope

Down-slope shape: Concave

Across-slope shape: Linear

Parent material: Volcanic ash over loess

Typical profile

Oi - 0 to 1 inches: slightly decomposed plant material

Oe - 1 to 2 inches: moderately decomposed plant material

A - 2 to 4 inches: ashy silt loam

BA - 4 to 15 inches: ashy silt loam

Bw - 15 to 22 inches: silt loam

Bt - 22 to 29 inches: silt loam

EBc - 29 to 31 inches: silt loam

Btxcb - 31 to 53 inches: silty clay loam

Btxb - 53 to 60 inches: silty clay loam

Properties and qualities

Slope: 3 to 8 percent

Depth to restrictive feature: 23 to 40 inches to fragipan

Natural drainage class: Moderately well drained

Capacity of the most limiting layer to transmit water (Ksat): Moderately high to high
(0.57 to 1.98 in/hr)

Depth to water table: About 16 to 22 inches

Frequency of flooding: None

Frequency of ponding: None

Available water storage in profile: Moderate (about 6.7 inches)

Interpretive groups

Land capability classification (irrigated): 3e

Land capability classification (nonirrigated): 3e

Hydrologic Soil Group: C/D

Other vegetative classification: Douglas-fir/ninebark (CN260)

Kootenai County Area, Idaho

1hn6j—Blinn ashy silt loam, 5 to 35 percent slopes, stony

Map Unit Setting

National map unit symbol: 1hn6j
Elevation: 2,100 to 3,000 feet
Mean annual precipitation: 25 to 30 inches
Mean annual air temperature: 43 to 45 degrees F
Frost-free period: 90 to 110 days
Farmland classification: Not prime farmland

Map Unit Composition

Blinn, stony surface, and similar soils: 80 percent
Estimates are based on observations, descriptions, and transects of the mapunit.

Description of Blinn, Stony Surface

Setting

Landform: Escarpments, structural benches
Landform position (two-dimensional): Backslope, shoulder
Landform position (three-dimensional): Side slope
Down-slope shape: Convex
Across-slope shape: Convex
Parent material: Volcanic ash and loess over colluvium over residuum weathered from basalt

Typical profile

Oi - 0 to 1 inches: slightly decomposed plant material
Oe - 1 to 2 inches: moderately decomposed plant material
A - 2 to 6 inches: ashy silt loam
Bw1 - 6 to 12 inches: gravelly ashy silt loam
Bw2 - 12 to 24 inches: stony loam
C - 24 to 39 inches: very stony loam
R - 39 to 40 inches: bedrock

Properties and qualities

Slope: 5 to 35 percent
Percent of area covered with surface fragments: 0.1 percent
Depth to restrictive feature: 20 to 40 inches to lithic bedrock
Natural drainage class: Well drained
Capacity of the most limiting layer to transmit water (Ksat): Moderately high to high (0.57 to 1.98 in/hr)
Depth to water table: More than 80 inches
Frequency of flooding: None
Frequency of ponding: None
Available water storage in profile: Low (about 5.1 inches)

Interpretive groups

Land capability classification (irrigated): None specified
Land capability classification (nonirrigated): 4e
Hydrologic Soil Group: C
Other vegetative classification: grand fir/ninebark (CN506)

135—Lacy gravelly loam, very stony-Rock outcrop complex, 5 to 35 percent slopes

Map Unit Setting

National map unit symbol: 2nn3
Elevation: 1,500 to 3,200 feet
Mean annual precipitation: 22 to 28 inches
Mean annual air temperature: 46 to 52 degrees F
Frost-free period: 100 to 140 days
Farmland classification: Not prime farmland

Map Unit Composition

Lacy, very stony surface, and similar soils: 55 percent
Rock outcrop: 35 percent
Estimates are based on observations, descriptions, and transects of the mapunit.

Description of Lacy, Very Stony Surface

Setting

Landform: Canyons, plateaus
Landform position (two-dimensional): Summit, shoulder
Down-slope shape: Convex
Across-slope shape: Linear
Parent material: Loess and/or colluvium over bedrock derived from basalt

Typical profile

Oi - 0 to 1 inches: slightly decomposed plant material
A - 1 to 8 inches: gravelly loam
Bt1 - 8 to 15 inches: stony clay loam
Bt2 - 15 to 19 inches: extremely stony clay loam
R - 19 to 30 inches: bedrock

Properties and qualities

Slope: 5 to 35 percent
Percent of area covered with surface fragments: 2.0 percent
Depth to restrictive feature: 10 to 20 inches to lithic bedrock
Natural drainage class: Well drained
Capacity of the most limiting layer to transmit water (Ksat): Moderately high to high (0.57 to 2.00 in/hr)
Depth to water table: More than 80 inches
Frequency of flooding: None
Frequency of ponding: None
Available water storage in profile: Very low (about 3.0 inches)

Interpretive groups

Land capability classification (irrigated): None specified
Land capability classification (nonirrigated): 6s
Hydrologic Soil Group: D
Other vegetative classification: ponderosa pine/Idaho fescue (CN140)

Description of Rock Outcrop

Typical profile

R - 0 to 60 inches: bedrock

Properties and qualities

Slope: 5 to 35 percent

Depth to restrictive feature: 0 inches to lithic bedrock

Interpretive groups

Land capability classification (irrigated): None specified

Land capability classification (nonirrigated): 8

146—McCrosket-Ardenvoir association, 20 to 35 percent slopes

Map Unit Setting

National map unit symbol: 2nng

Elevation: 2,200 to 5,000 feet

Mean annual precipitation: 24 to 35 inches

Mean annual air temperature: 43 to 50 degrees F

Frost-free period: 90 to 120 days

Farmland classification: Not prime farmland

Map Unit Composition

Mccrosket and similar soils: 55 percent

Ardenvoir and similar soils: 35 percent

Estimates are based on observations, descriptions, and transects of the mapunit.

Description of Mccrosket

Setting

Landform: Ridges, mountains

Landform position (two-dimensional): Shoulder, backslope

Down-slope shape: Convex

Across-slope shape: Convex

Parent material: Volcanic ash and/or loess over residuum weathered from metasedimentary rock

Typical profile

Oi - 0 to 1 inches: slightly decomposed plant material

Oe - 1 to 2 inches: moderately decomposed plant material

A - 2 to 10 inches: gravelly silt loam

Bw - 10 to 48 inches: extremely cobbly silt loam

Cr - 48 to 58 inches: bedrock

Properties and qualities

Slope: 20 to 35 percent

Depth to restrictive feature: 40 to 60 inches to paralithic bedrock

Natural drainage class: Well drained

Capacity of the most limiting layer to transmit water (Ksat): Moderately high to high
(0.57 to 2.00 in/hr)

Custom Soil Resource Report

Depth to water table: More than 80 inches
Frequency of flooding: None
Frequency of ponding: None
Available water storage in profile: Low (about 5.2 inches)

Interpretive groups

Land capability classification (irrigated): None specified
Land capability classification (nonirrigated): 6e
Hydrologic Soil Group: B

Description of Ardenvoir

Setting

Landform: Mountains
Landform position (two-dimensional): Backslope
Down-slope shape: Concave
Across-slope shape: Linear
Parent material: Volcanic ash and loess over residuum weathered from metasedimentary rock

Typical profile

Oi - 0 to 1 inches: slightly decomposed plant material
AB - 1 to 17 inches: gravelly loam
BC - 17 to 48 inches: very cobbly loam
Cr - 48 to 58 inches: bedrock

Properties and qualities

Slope: 20 to 35 percent
Depth to restrictive feature: 40 to 60 inches to paralithic bedrock
Natural drainage class: Well drained
Capacity of the most limiting layer to transmit water (Ksat): Moderately high to high (0.57 to 2.00 in/hr)
Depth to water table: More than 80 inches
Frequency of flooding: None
Frequency of ponding: None
Available water storage in profile: Low (about 5.6 inches)

Interpretive groups

Land capability classification (irrigated): None specified
Land capability classification (nonirrigated): 6e
Hydrologic Soil Group: B
Other vegetative classification: grand fir/twinflower (CN590)

157—Porrett silt loam

Map Unit Setting

National map unit symbol: 2nnt
Elevation: 2,100 to 2,900 feet
Mean annual precipitation: 25 to 35 inches
Mean annual air temperature: 41 to 45 degrees F
Frost-free period: 60 to 90 days

Custom Soil Resource Report

Farmland classification: Prime farmland if drained and either protected from flooding or not frequently flooded during the growing season

Map Unit Composition

Porrett and similar soils: 85 percent

Minor components: 5 percent

Estimates are based on observations, descriptions, and transects of the mapunit.

Description of Porrett

Setting

Landform: Drainageways, stream terraces

Down-slope shape: Concave

Across-slope shape: Linear

Parent material: Volcanic ash and alluvium and loess

Typical profile

Ap - 0 to 3 inches: ashy silt loam

E - 3 to 28 inches: silt loam

Btg - 28 to 60 inches: silty clay loam

Properties and qualities

Slope: 0 to 2 percent

Depth to restrictive feature: More than 80 inches

Natural drainage class: Very poorly drained

Capacity of the most limiting layer to transmit water (Ksat): Moderately high (0.20 to 0.60 in/hr)

Depth to water table: About 0 to 12 inches

Frequency of flooding: Frequent

Frequency of ponding: None

Available water storage in profile: High (about 12.0 inches)

Interpretive groups

Land capability classification (irrigated): 6e

Land capability classification (nonirrigated): 5w

Hydrologic Soil Group: C/D

Minor Components

Potlatch

Percent of map unit: 5 percent

Landform: Flood plains

165—Santa ashy silt loam, 2 to 8 percent slopes

Map Unit Setting

National map unit symbol: 2v71x

Elevation: 2,610 to 3,200 feet

Mean annual precipitation: 25 to 35 inches

Mean annual air temperature: 43 to 46 degrees F

Frost-free period: 110 to 120 days

Farmland classification: Farmland of statewide importance, if drained

Map Unit Composition

Santa and similar soils: 85 percent

Estimates are based on observations, descriptions, and transects of the mapunit.

Description of Santa

Setting

Landform: Interfluves, hillslopes

Landform position (two-dimensional): Summit, shoulder, backslope

Landform position (three-dimensional): Interfluve, side slope

Down-slope shape: Convex, concave

Across-slope shape: Linear, convex

Parent material: Mixed volcanic ash and loess

Typical profile

Ap - 0 to 8 inches: ashy silt loam

Bw - 8 to 19 inches: silt loam

E - 19 to 29 inches: silt loam

Btxb1 - 29 to 38 inches: silt loam

Btxb2 - 38 to 59 inches: silt loam

Properties and qualities

Slope: 2 to 8 percent

Depth to restrictive feature: 24 to 39 inches to fragipan

Natural drainage class: Moderately well drained

Capacity of the most limiting layer to transmit water (Ksat): Very low to moderately low (0.00 to 0.06 in/hr)

Depth to water table: About 8 to 22 inches

Frequency of flooding: None

Frequency of ponding: None

Available water storage in profile: Low (about 5.7 inches)

Interpretive groups

Land capability classification (irrigated): None specified

Land capability classification (nonirrigated): 4e

Hydrologic Soil Group: C/D

Other vegetative classification: grand fir/ninebark (CN506)

166—Santa ashy silt loam, 8 to 15 percent slopes

Map Unit Setting

National map unit symbol: 2v71y

Elevation: 2,510 to 3,200 feet

Mean annual precipitation: 25 to 37 inches

Mean annual air temperature: 43 to 46 degrees F

Frost-free period: 105 to 120 days

Farmland classification: Not prime farmland

Map Unit Composition

Santa and similar soils: 80 percent

Estimates are based on observations, descriptions, and transects of the mapunit.

Description of Santa

Setting

Landform: Interfluves, hillslopes

Landform position (two-dimensional): Summit, shoulder, backslope

Landform position (three-dimensional): Interfluve, side slope

Down-slope shape: Convex, concave

Across-slope shape: Linear, convex

Parent material: Mixed volcanic ash and loess

Typical profile

Ap - 0 to 8 inches: ashy silt loam

Bw - 8 to 19 inches: silt loam

E - 19 to 29 inches: silt loam

Btxb1 - 29 to 38 inches: silt loam

Btxb2 - 38 to 59 inches: silt loam

Properties and qualities

Slope: 8 to 15 percent

Depth to restrictive feature: 24 to 39 inches to fragipan

Natural drainage class: Moderately well drained

Capacity of the most limiting layer to transmit water (Ksat): Very low to moderately low (0.00 to 0.06 in/hr)

Depth to water table: About 8 to 22 inches

Frequency of flooding: None

Frequency of ponding: None

Available water storage in profile: Low (about 5.7 inches)

Interpretive groups

Land capability classification (irrigated): None specified

Land capability classification (nonirrigated): 4e

Hydrologic Soil Group: C/D

Other vegetative classification: grand fir/ninebark (CN506)

168—Santa variant silt loam, 5 to 20 percent slopes

Map Unit Setting

National map unit symbol: 2np5

Elevation: 2,300 to 3,000 feet

Mean annual precipitation: 26 to 28 inches

Mean annual air temperature: 43 to 45 degrees F

Frost-free period: 100 to 120 days

Farmland classification: Not prime farmland

Map Unit Composition

Santa variant and similar soils: 80 percent

Estimates are based on observations, descriptions, and transects of the mapunit.

Description of Santa Variant

Setting

Landform: Hillslopes

Down-slope shape: Convex

Across-slope shape: Linear

Parent material: Loess over bedrock derived from basalt and/or metasedimentary rock

Typical profile

Oi - 0 to 1 inches: slightly decomposed plant material

AB - 1 to 10 inches: silt loam

E - 10 to 24 inches: silt loam

Btx - 24 to 37 inches: silty clay loam

R - 37 to 47 inches: bedrock

Properties and qualities

Slope: 5 to 20 percent

Depth to restrictive feature: 20 to 40 inches to lithic bedrock; 20 to 36 inches to fragipan

Natural drainage class: Moderately well drained

Capacity of the most limiting layer to transmit water (Ksat): Very low to moderately low (0.00 to 0.06 in/hr)

Depth to water table: About 18 to 30 inches

Frequency of flooding: None

Frequency of ponding: None

Available water storage in profile: Low (about 5.1 inches)

Interpretive groups

Land capability classification (irrigated): None specified

Land capability classification (nonirrigated): 4e

Hydrologic Soil Group: C

Other vegetative classification: grand fir/twinflower (CN590)

186—Taney silt loam, 3 to 7 percent slopes

Map Unit Setting

National map unit symbol: 2npr

Elevation: 2,300 to 3,900 feet

Mean annual precipitation: 24 to 27 inches

Mean annual air temperature: 43 to 45 degrees F

Frost-free period: 100 to 130 days

Farmland classification: Prime farmland if drained

Map Unit Composition

Taney and similar soils: 75 percent

Estimates are based on observations, descriptions, and transects of the mapunit.

Description of Taney

Setting

Landform: Hillslopes

Custom Soil Resource Report

Landform position (two-dimensional): Summit, backslope
Down-slope shape: Convex
Across-slope shape: Linear
Parent material: Volcanic ash and/or loess

Typical profile

A - 0 to 21 inches: silt loam
E - 21 to 26 inches: silt loam
Btx - 26 to 60 inches: silty clay loam

Properties and qualities

Slope: 3 to 7 percent
Depth to restrictive feature: 26 to 40 inches to fragipan
Natural drainage class: Moderately well drained
Capacity of the most limiting layer to transmit water (Ksat): Very low to moderately low (0.00 to 0.06 in/hr)
Depth to water table: About 18 to 30 inches
Frequency of flooding: None
Frequency of ponding: None
Available water storage in profile: Low (about 5.7 inches)

Interpretive groups

Land capability classification (irrigated): 4s
Land capability classification (nonirrigated): 4s
Hydrologic Soil Group: C
Other vegetative classification: Douglas-fir/common snowberry (CN310)

187—Taney silt loam, 7 to 25 percent slopes

Map Unit Setting

National map unit symbol: 2nps
Elevation: 2,300 to 3,900 feet
Mean annual precipitation: 24 to 27 inches
Mean annual air temperature: 43 to 45 degrees F
Frost-free period: 100 to 130 days
Farmland classification: Farmland of statewide importance, if drained

Map Unit Composition

Taney and similar soils: 75 percent
Estimates are based on observations, descriptions, and transects of the mapunit.

Description of Taney

Setting

Landform: Hillslopes
Landform position (two-dimensional): Summit, backslope
Down-slope shape: Linear
Across-slope shape: Concave
Parent material: Volcanic ash and/or loess

Typical profile

A - 0 to 21 inches: silt loam
E - 21 to 26 inches: silt loam

Btx - 26 to 60 inches: silty clay loam

Properties and qualities

Slope: 7 to 25 percent

Depth to restrictive feature: 26 to 40 inches to fragipan

Natural drainage class: Moderately well drained

Capacity of the most limiting layer to transmit water (Ksat): Very low to moderately low (0.00 to 0.06 in/hr)

Depth to water table: About 18 to 30 inches

Frequency of flooding: None

Frequency of ponding: None

Available water storage in profile: Low (about 5.7 inches)

Interpretive groups

Land capability classification (irrigated): None specified

Land capability classification (nonirrigated): 4e

Hydrologic Soil Group: C

Other vegetative classification: Douglas-fir/common snowberry (CN310)

189—Tekoa gravelly silt loam, 5 to 20 percent slopes

Map Unit Setting

National map unit symbol: 2npv

Elevation: 2,000 to 4,000 feet

Mean annual precipitation: 20 to 30 inches

Mean annual air temperature: 45 to 48 degrees F

Frost-free period: 100 to 140 days

Farmland classification: Farmland of statewide importance

Map Unit Composition

Tekoa and similar soils: 80 percent

Estimates are based on observations, descriptions, and transects of the mapunit.

Description of Tekoa

Setting

Landform: Mountains

Landform position (two-dimensional): Backslope

Down-slope shape: Convex

Across-slope shape: Convex

Parent material: Volcanic ash and loess over residuum weathered from quartzite

Typical profile

Oi - 0 to 1 inches: slightly decomposed plant material

Oe - 1 to 2 inches: moderately decomposed plant material

A - 2 to 9 inches: gravelly silt loam

Bt - 9 to 16 inches: very gravelly silt loam

BC - 16 to 32 inches: very gravelly silt loam

Cr - 32 to 42 inches: bedrock

Properties and qualities

Slope: 5 to 20 percent

Depth to restrictive feature: 20 to 40 inches to paralithic bedrock

Custom Soil Resource Report

Natural drainage class: Well drained

Capacity of the most limiting layer to transmit water (Ksat): Moderately high to high
(0.57 to 2.00 in/hr)

Depth to water table: More than 80 inches

Frequency of flooding: None

Frequency of ponding: None

Available water storage in profile: Low (about 4.0 inches)

Interpretive groups

Land capability classification (irrigated): None specified

Land capability classification (nonirrigated): 4e

Hydrologic Soil Group: C

Other vegetative classification: ponderosa pine/Idaho fescue (CN140)

232—Lacy-Bobbitt complex, 5 to 35 percent slopes, stony

Map Unit Setting

National map unit symbol: 1hn6s

Elevation: 2,130 to 3,000 feet

Mean annual precipitation: 25 to 30 inches

Mean annual air temperature: 47 to 50 degrees F

Frost-free period: 100 to 130 days

Farmland classification: Not prime farmland

Map Unit Composition

Lacy, stony surface, and similar soils: 55 percent

Bobbitt, stony surface, and similar soils: 30 percent

Estimates are based on observations, descriptions, and transects of the mapunit.

Description of Lacy, Stony Surface

Setting

Landform: Escarpments, structural benches

Landform position (two-dimensional): Backslope, shoulder

Landform position (three-dimensional): Side slope

Down-slope shape: Convex

Across-slope shape: Convex

Parent material: Loess over colluvium over bedrock derived from basalt

Typical profile

Oi - 0 to 1 inches: slightly decomposed plant material

Oe - 1 to 2 inches: moderately decomposed plant material

A1 - 2 to 3 inches: stony silt loam

A2 - 3 to 10 inches: stony silt loam

Bt1 - 10 to 14 inches: very stony silt loam

Bt2 - 14 to 17 inches: extremely stony loam

R - 17 to 27 inches: bedrock

Properties and qualities

Slope: 5 to 35 percent

Percent of area covered with surface fragments: 0.1 percent

Depth to restrictive feature: 10 to 20 inches to lithic bedrock

Custom Soil Resource Report

Natural drainage class: Well drained

Capacity of the most limiting layer to transmit water (Ksat): Moderately high to high
(0.57 to 1.98 in/hr)

Depth to water table: More than 80 inches

Frequency of flooding: None

Frequency of ponding: None

Available water storage in profile: Very low (about 2.5 inches)

Interpretive groups

Land capability classification (irrigated): None specified

Land capability classification (nonirrigated): 6e

Hydrologic Soil Group: D

Other vegetative classification: ponderosa pine/common snowberry (CN170)

Description of Bobbitt, Stony Surface

Setting

Landform: Escarpments, structural benches

Landform position (two-dimensional): Backslope, shoulder

Landform position (three-dimensional): Side slope

Down-slope shape: Convex

Across-slope shape: Convex

Parent material: Volcanic ash and loess over colluvium over residuum weathered from basalt

Typical profile

Oi - 0 to 1 inches: slightly decomposed plant material

Oe - 1 to 2 inches: moderately decomposed plant material

A - 2 to 9 inches: stony ashy silt loam

Bt - 9 to 23 inches: very stony clay loam

R - 23 to 33 inches: bedrock

Properties and qualities

Slope: 5 to 35 percent

Percent of area covered with surface fragments: 0.1 percent

Depth to restrictive feature: 20 to 40 inches to lithic bedrock

Natural drainage class: Well drained

Capacity of the most limiting layer to transmit water (Ksat): Moderately high to high
(0.57 to 1.98 in/hr)

Depth to water table: More than 80 inches

Frequency of flooding: None

Frequency of ponding: None

Available water storage in profile: Low (about 3.3 inches)

Interpretive groups

Land capability classification (irrigated): None specified

Land capability classification (nonirrigated): 4e

Hydrologic Soil Group: C

Other vegetative classification: Douglas-fir/ninebark (CN260)

335—Carlinton ashy silt loam, dry, 8 to 25 percent slopes

Map Unit Setting

National map unit symbol: 1hn7g

Elevation: 2,700 to 3,230 feet

Mean annual precipitation: 25 to 28 inches

Mean annual air temperature: 43 to 45 degrees F

Frost-free period: 90 to 120 days

Farmland classification: Farmland of statewide importance

Map Unit Composition

Carlinton, dry, and similar soils: 80 percent

Minor components: 2 percent

Estimates are based on observations, descriptions, and transects of the mapunit.

Description of Carlinton, Dry

Setting

Landform: Loess hills

Landform position (two-dimensional): Backslope

Landform position (three-dimensional): Side slope

Down-slope shape: Convex

Across-slope shape: Linear

Parent material: Volcanic ash over loess

Typical profile

Ap1 - 0 to 5 inches: ashy silt loam

Ap2 - 5 to 10 inches: ashy silt loam

Bw - 10 to 14 inches: silt loam

EBt - 14 to 20 inches: silt loam

E - 20 to 23 inches: silt loam

BtbxE - 23 to 30 inches: silt loam

Btbx - 30 to 53 inches: silty clay loam

Btb - 53 to 60 inches: silty clay loam

Properties and qualities

Slope: 8 to 25 percent

Depth to restrictive feature: 21 to 40 inches to fragipan

Natural drainage class: Moderately well drained

Capacity of the most limiting layer to transmit water (Ksat): Moderately high to high
(0.57 to 1.98 in/hr)

Depth to water table: About 14 to 20 inches

Frequency of flooding: None

Frequency of ponding: None

Available water storage in profile: Low (about 5.7 inches)

Interpretive groups

Land capability classification (irrigated): None specified

Land capability classification (nonirrigated): 3e

Hydrologic Soil Group: C/D

Other vegetative classification: Douglas-fir/ninebark (CN260)

Minor Components

Lovell

Percent of map unit: 2 percent
Landform: Drainageways
Landform position (two-dimensional): Toeslope
Down-slope shape: Concave
Across-slope shape: Linear
Ecological site: MEADOW (R009XY018ID)

336—Carlinton, dry-Taney complex, 3 to 8 percent slopes

Map Unit Setting

National map unit symbol: 1hn7h
Elevation: 2,560 to 3,020 feet
Mean annual precipitation: 25 to 28 inches
Mean annual air temperature: 43 to 45 degrees F
Frost-free period: 90 to 120 days
Farmland classification: All areas are prime farmland

Map Unit Composition

Carlinton, dry, and similar soils: 55 percent
Taney and similar soils: 25 percent
Estimates are based on observations, descriptions, and transects of the mapunit.

Description of Carlinton, Dry

Setting

Landform: Loess hills
Landform position (two-dimensional): Shoulder
Landform position (three-dimensional): Side slope
Down-slope shape: Convex
Across-slope shape: Linear
Parent material: Volcanic ash over loess

Typical profile

Ap1 - 0 to 5 inches: ashy silt loam
Ap2 - 5 to 10 inches: ashy silt loam
Bw - 10 to 14 inches: silt loam
EBt - 14 to 20 inches: silt loam
E - 20 to 23 inches: silt loam
BtbxE - 23 to 30 inches: silt loam
Btbx - 30 to 53 inches: silty clay loam
Btb - 53 to 60 inches: silty clay loam

Properties and qualities

Slope: 3 to 8 percent
Depth to restrictive feature: 21 to 40 inches to fragipan
Natural drainage class: Moderately well drained

Custom Soil Resource Report

Capacity of the most limiting layer to transmit water (Ksat): Moderately high to high
(0.57 to 1.98 in/hr)

Depth to water table: About 14 to 20 inches

Frequency of flooding: None

Frequency of ponding: None

Available water storage in profile: Low (about 5.7 inches)

Interpretive groups

Land capability classification (irrigated): 3e

Land capability classification (nonirrigated): 3e

Hydrologic Soil Group: C/D

Other vegetative classification: Douglas-fir/ninebark (CN260)

Description of Taney

Setting

Landform: Loess hills

Landform position (two-dimensional): Shoulder, summit

Landform position (three-dimensional): Side slope

Down-slope shape: Concave

Across-slope shape: Linear

Parent material: Volcanic ash over loess

Typical profile

Oi - 0 to 1 inches: slightly decomposed plant material

Oe - 1 to 2 inches: moderately decomposed plant material

A - 2 to 4 inches: ashy silt loam

BA - 4 to 15 inches: ashy silt loam

Bw - 15 to 22 inches: silt loam

Bt - 22 to 29 inches: silt loam

EBc - 29 to 31 inches: silt loam

Btxcb - 31 to 53 inches: silty clay loam

Btxb - 53 to 60 inches: silty clay loam

Properties and qualities

Slope: 3 to 8 percent

Depth to restrictive feature: 23 to 40 inches to fragipan

Natural drainage class: Moderately well drained

Capacity of the most limiting layer to transmit water (Ksat): Moderately high to high
(0.57 to 1.98 in/hr)

Depth to water table: About 16 to 22 inches

Frequency of flooding: None

Frequency of ponding: None

Available water storage in profile: Moderate (about 6.7 inches)

Interpretive groups

Land capability classification (irrigated): 3e

Land capability classification (nonirrigated): 3e

Hydrologic Soil Group: C/D

Other vegetative classification: Douglas-fir/ninebark (CN260)

References

American Association of State Highway and Transportation Officials (AASHTO). 2004. Standard specifications for transportation materials and methods of sampling and testing. 24th edition.

American Society for Testing and Materials (ASTM). 2005. Standard classification of soils for engineering purposes. ASTM Standard D2487-00.

Cowardin, L.M., V. Carter, F.C. Golet, and E.T. LaRoe. 1979. Classification of wetlands and deep-water habitats of the United States. U.S. Fish and Wildlife Service FWS/OBS-79/31.

Federal Register. July 13, 1994. Changes in hydric soils of the United States.

Federal Register. September 18, 2002. Hydric soils of the United States.

Hurt, G.W., and L.M. Vasilas, editors. Version 6.0, 2006. Field indicators of hydric soils in the United States.

National Research Council. 1995. Wetlands: Characteristics and boundaries.

Soil Survey Division Staff. 1993. Soil survey manual. Soil Conservation Service. U.S. Department of Agriculture Handbook 18. http://www.nrcs.usda.gov/wps/portal/nrcs/detail/national/soils/?cid=nrcs142p2_054262

Soil Survey Staff. 1999. Soil taxonomy: A basic system of soil classification for making and interpreting soil surveys. 2nd edition. Natural Resources Conservation Service, U.S. Department of Agriculture Handbook 436. http://www.nrcs.usda.gov/wps/portal/nrcs/detail/national/soils/?cid=nrcs142p2_053577

Soil Survey Staff. 2010. Keys to soil taxonomy. 11th edition. U.S. Department of Agriculture, Natural Resources Conservation Service. http://www.nrcs.usda.gov/wps/portal/nrcs/detail/national/soils/?cid=nrcs142p2_053580

Tiner, R.W., Jr. 1985. Wetlands of Delaware. U.S. Fish and Wildlife Service and Delaware Department of Natural Resources and Environmental Control, Wetlands Section.

United States Army Corps of Engineers, Environmental Laboratory. 1987. Corps of Engineers wetlands delineation manual. Waterways Experiment Station Technical Report Y-87-1.

United States Department of Agriculture, Natural Resources Conservation Service. National forestry manual. http://www.nrcs.usda.gov/wps/portal/nrcs/detail/soils/home/?cid=nrcs142p2_053374

United States Department of Agriculture, Natural Resources Conservation Service. National range and pasture handbook. <http://www.nrcs.usda.gov/wps/portal/nrcs/detail/national/landuse/rangepasture/?cid=stelprdb1043084>

Custom Soil Resource Report

United States Department of Agriculture, Natural Resources Conservation Service. National soil survey handbook, title 430-VI. http://www.nrcs.usda.gov/wps/portal/nrcs/detail/soils/scientists/?cid=nrcs142p2_054242

United States Department of Agriculture, Natural Resources Conservation Service. 2006. Land resource regions and major land resource areas of the United States, the Caribbean, and the Pacific Basin. U.S. Department of Agriculture Handbook 296. http://www.nrcs.usda.gov/wps/portal/nrcs/detail/national/soils/?cid=nrcs142p2_053624

United States Department of Agriculture, Soil Conservation Service. 1961. Land capability classification. U.S. Department of Agriculture Handbook 210. http://www.nrcs.usda.gov/Internet/FSE_DOCUMENTS/nrcs142p2_052290.pdf