4° 41' 45" W

30° 40' 26" N

30° 40' 26" N



30° 39′ 50″ N

N

Map Scale: 1:7,920 if printed on A landscape (11"  $\times$  8.5") sheet.

0 100 200 400 600 Feet
0 350 700 1400 2100
Map projection: Web Mercator Corner coordinates: WGS84



30° 39' 50" N

#### MAP LEGEND MAP INFORMATION The soil surveys that comprise your AOI were mapped at Area of Interest (AOI) Coarse-loamy, siliceous, Fine-loamy, kaolinitic, thermic Typic active, acid, thermic Typic 1:12.000. Area of Interest (AOI) Fluvaquents Kanhapludults Please rely on the bar scale on each map sheet for map Soils Fine-loamy, kaolinitic, Loamy, kaolinitic, thermic measurements. thermic Plinthic Arenic Plinthic Soil Rating Polygons Kandiudults Kandiudults Coarse-loamy, siliceous, Source of Map: Natural Resources Conservation Service Fine-loamy, kaolinitic, Loamy, siliceous, active, acid, thermic Typic Web Soil Survey URL: http://websoilsurvey.nrcs.usda.gov thermic Typic subactive, thermic Arenic Fluvaquents Coordinate System: Web Mercator (EPSG:3857) Kanhapludults Plinthaquic Paleudults Fine-loamy, kaolinitic, Loamy, kaolinitic, thermic Loamy, siliceous, Maps from the Web Soil Survey are based on the Web Mercator thermic Plinthic Arenic Plinthic subactive, thermic Kandiudults projection, which preserves direction and shape but distorts Grossarenic Plinthic Kandiudults distance and area. A projection that preserves area, such as the Fine-loamy, kaolinitic, Paleudults Loamy, siliceous, thermic Typic Albers equal-area conic projection, should be used if more subactive, thermic Arenic Udorthents Kanhapludults accurate calculations of distance or area are required. Plinthaguic Paleudults Loamy, kaolinitic, thermic Not rated or not available This product is generated from the USDA-NRCS certified data as Loamy, siliceous, Arenic Plinthic of the version date(s) listed below. subactive, thermic Kandiudults **Water Features** Grossarenic Plinthic Loamy, siliceous, Streams and Canals Soil Survey Area: Gadsden County, Florida Paleudults subactive, thermic Arenic Survey Area Data: Version 19, Dec 27, 2013 Udorthents Transportation Plinthaguic Paleudults Rails Soil map units are labeled (as space allows) for map scales Loamy, siliceous, Not rated or not available subactive, thermic 1:50.000 or larger. Interstate Highways Grossarenic Plinthic Soil Rating Points Paleudults Date(s) aerial images were photographed: Apr 4, 2010—Jan Coarse-loamy, siliceous, **US Routes** 12, 2011 Udorthents active, acid, thermic Typic Major Roads Fluvaquents Not rated or not available The orthophoto or other base map on which the soil lines were Fine-loamy, kaolinitic, Local Roads compiled and digitized probably differs from the background $\sim$ thermic Plinthic Soil Rating Lines imagery displayed on these maps. As a result, some minor shifting Kandiudults Background of map unit boundaries may be evident. Aerial Photography

# **Soil Taxonomy Classification**

Soil Tax	Soil Taxonomy Classification— Summary by Map Unit — Gadsden County, Florida (FL039)	Summary by Map Unit —	Gadsden County, Florida	(FL039)
Map unit symbol	Map unit name	Rating	Acres in AOI	Percent of AOI
14	Cowarts-Dothan-Fuquay complex, 5 to 8 percent slopes	Fine-loamy, kaolinitic, thermic Typic Kanhapludults	51.5	35.6%
19	Dothan-Fuquay complex, 2 to 5 percent slopes	Fine-loamy, kaolinitic, thermic Plinthic Kandiudults	5.9	4.1%
21	Dothan-Fuquay-Cowarts complex, 8 to 15 percent slopes	Fine-loamy, kaolinitic, thermic Plinthic Kandiudults	10.4	7.2%
24	Fuquay-Bonifay complex, 5 to 15 percent slopes	Loamy, kaolinitic, thermic Arenic Plinthic Kandiudults	6.0	4.2%
65	Udorthents, reclaimed	Udorthents	5.1	3.5%
77	Bonifay-Fuquay complex, 0 to 5 percent slopes	Loamy, siliceous, subactive, thermic Grossarenic Plinthic Paleudults	10.7	7.4%
89	Bibb-Rains-Garcon complex, occasionally flooded	Coarse-loamy, siliceous, active, acid, thermic Typic Fluvaquents	9.8	6.8%
99	Water		1.7	1.2%
107	Fuquay-Bonifay complex, 0 to 5 percent slopes	Loamy, kaolinitic, thermic Arenic Plinthic Kandiudults	42.9	29.7%
113	Leefield fine sand, 0 to 5 percent slopes	Loamy, siliceous, subactive, thermic Arenic Plinthaquic Paleudults	0.6	0.4%
Totals for Area of Interest	est		144.7	100.0%

### **Description**

This rating presents the taxonomic classification based on Soil Taxonomy

Classification is based on soil properties observed in the field or inferred from those observations or from laboratory measurements. This table shows the classification The system of soil classification used by the National Cooperative Soil Survey has six categories (Soil Survey Staff, 1999 and 2003). Beginning with the broadest, of the soils in the survey area. The categories are defined in the following these categories are the order, suborder, great group, subgroup, family, and series

is identified by a word ending in sol. An example is Alfisols. the dominant soil-forming processes and the degree of soil formation. Each order ORDER. Twelve soil orders are recognized. The differences among orders reflect

name of a suborder indicates the order. An example is Udalfs (Ud, meaning humid plus alfs, from Alfisols). that reflect the most important variables within the orders. The last syllable in the properties that influence soil genesis and are important to plant growth or properties SUBORDER. Each order is divided into suborders primarily on the basis of

horizonation, plus udalfs, the suborder of the Alfisols that has a udic moisture indicates a property of the soil. An example is Hapludalfs (Hapl, meaning minimal status. Each great group is identified by the name of a suborder and by a prefix that horizons; soil moisture and temperature regimes; type of saturation; and base similarities in kind, arrangement, and degree of development of pedogenic GREAT GROUP. Each suborder is divided into great groups on the basis of close

orders, suborders, or great groups. Extragrades have some properties that are not group; it is not necessarily the most extensive. Intergrades are transitions to other the great group. An example is Typic Hapludalfs. the name of the great group. The adjective Typic identifies the subgroup that typifies taxonomic class. Each subgroup is identified by one or more adjectives preceding representative of the great group but do not indicate transitions to any other intergrades or extragrades. The typic subgroup is the central concept of the great SUBGROUP. Each great group has a typic subgroup. Other subgroups are

fine-loamy, mixed, active, mesic Typic Hapludalfs. name of a subgroup preceded by terms that indicate soil properties. An example is temperature regime, soil depth, and reaction class. A family name consists of the particle-size class, mineralogy class, cation-exchange activity class, soil biological activity. Among the properties and characteristics considered are the properties are those of horizons below plow depth where there is much chemical properties and other characteristics that affect management. Generally, FAMILY. Families are established within a subgroup on the basis of physical and

and arrangement in the profile. color, texture, structure, reaction, consistence, mineral and chemical composition, SERIES. The series consists of soils within a family that have horizons similar in



#### References:

Soil Survey Staff. 1999. Soil taxonomy: A basic system of soil classification for making and interpreting soil surveys. 2nd edition. Natural Resources Conservation Service. U.S. Department of Agriculture Handbook 436.

Soil Survey Staff. 2006. Keys to soil taxonomy. 10th edition. U.S. Department of Agriculture, Natural Resources Conservation Service. (The soils in a given survey area may have been classified according to earlier editions of this publication.)

## **Rating Options**

Aggregation Method: Dominant Condition

Component Percent Cutoff: None Specified

Tie-break Rule: Lower