Ironwood 32

Iowa, 32 AC +/-





| Boundary 32.5 ac

| SOIL CODE | SOIL DESCRIPTION | ACRES | % | CSR2 | CAP |
|-----------|---|-------|-------|-------|-----|
| LcD2 | Lamoni clay loam, 9 to 14 percent slopes, moderately eroded | 0.7 | 2.28 | 10 | 4e |
| LaD2 | Ladoga silt loam, 9 to 14 percent slopes, eroded | 0.0 | 0.08 | 49 | 3e |
| SbC2 | Sharpsburg silty clay loam, 5 to 9 percent slopes, eroded | 3.9 | 12.13 | 80 | 3e |
| LaB | Ladoga silt loam, 2 to 5 percent slopes | 2.8 | 8.63 | 86 | 2e |
| LaC2 | Ladoga silt loam, dissected till plain, 5 to 9 percent slopes, eroded | 10.1 | 31.0 | 75 | 3e |
| SIE2 | Shelby-Lamoni complex, 14 to 18 percent slopes, moderately eroded | 0.2 | 0.47 | 27 | 6e |
| SbB | Sharpsburg silty clay loam, 2 to 5 percent slopes | 14.8 | 45.41 | 91 | 2e |
| TOTALS | | 32.5 | 100% | 82.09 | 2.5 |

| Capability Legend | | | | | | | | | | | | |
|--|---|---|---|---|---|---|---|---|--|--|--|--|
| Increased Limitations and Hazards | | | | | | | | | | | | |
| Decreased Adaptability and Freedom of Choice Users | | | | | | | | | | | | |
| Land, Capability Class(non-irrigated) | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | | | | |
| 'Wild Life' | • | • | • | • | • | • | • | • | | | | |
| Forestry | • | • | • | • | • | • | • | | | | | |
| Limited | • | • | • | • | • | • | • | | | | | |
| Moderate | • | • | • | • | • | • | | | | | | |
| Intense | • | • | • | • | • | | | | | | | |
| Limited | • | • | • | • | | | | | | | | |
| Moderate | • | • | • | | | | | | | | | |
| Intense | • | • | | | | | | | | | | |
| Very Intense | • | | | | | | | | | | | |
| | | | | | | | | | | | | |

Grazing Cultivation

- (c) climatic limitations (e) susceptibility to erosion
- (s) soil limitations within the rooting zone (w) excess of water