

# Herminjard Place 320 +/- ac. Panola County, Miss.

**TOTAL SIZE** 320+/- tax acres.

207.7+/- acres CRP (pines); 18.6+/- FSA cropland acres; 80+/- hardwood acres

17+/- acres of roads, ditches and clearings

**LOCATION** 7.5 miles northeast of Como, Panola County, MS; 22 miles north of Batesville, MS,

and 39 miles south of Memphis, TN. In Section 22, Twp 6S, Range 6W.

**DESCRIPTION** This is a prime deer and turkey hunting property and is accessible in all weather

by paved county roads: Paul White Road and Tom Floyd Road. The acres enrolled in CRP were planted to pine in 1990 and 1992. The woods and pine timber provide excellent habitat for deer and turkey, which are plentiful throughout the area. Trails have been established through the timber providing easy access to all hunting locations. The property is conveniently located just 9 miles from Sardis Reservoir for fishing and boating opportunities. The 18.6+/- acres of excellent bottom land cropland is leased to a neighboring farm. Future income will be generated through

the sale of pine timber.

**SOILS** Predominant cropland soil types are Grenada and Collins silt loam.

**PROPERTY TAXES** 2017 - \$2,141.04 or \$6.69 per acre.

**POSSESSION** At closing, subject to cropland lease.

**SHOWING** By appointment only.

**MANAGEMENT** Professional farm management available.






**CONTACT:** Murry McClintock, Broker, McClintock Land Company

Office: 662-363-1801 Cell: 662-363-6024 Fax: 662-363-1800


E-mail: murry@mcclintockland.com Address: PO Box 115, Tunica, MS 38676

## Herminjard Place (320 +/- ac.) Panola Co., Miss.

#### **Soils Details**








| Area Symbol: MS107, Soil Area Version: 12 | *c:_Using Capabilities Class Dominant Condition Aggregation Method |
|-------------------------------------------|--------------------------------------------------------------------|


|                                                        | Soil Description                                                                |       |          | Non-Irr         | Non-               | Bahiagrass | Common       | Corn  | Cotton | Grass         | Improved     | Pasture | Soybeans | Tall   | Tobacco |
|--------------------------------------------------------|---------------------------------------------------------------------------------|-------|----------|-----------------|--------------------|------------|--------------|-------|--------|---------------|--------------|---------|----------|--------|---------|
|                                                        |                                                                                 |       | of field | Class<br>Legend | Irr *c<br>Class *c |            | bermudagrass |       | lint   | legume<br>hay | bermudagrass |         |          | fescue |         |
| Gu                                                     | Gullied land,<br>silty                                                          | 92.73 | 29.0%    |                 | VIIe               |            |              |       |        |               |              |         |          |        |         |
| GrC3                                                   | Grenada silt<br>loam, 5 to 8<br>percent slopes,<br>severely<br>eroded           | 51.65 | 16.1%    |                 | ľVe                |            |              | 55    | 550    | 3.5           | 7            |         | 30       | 7      |         |
| Co                                                     | Collins silt<br>loam, local<br>alluvium<br>(occasionally<br>flooded)            | 46.66 | 14.6%    |                 | llw                |            |              | 110   | 800    |               | 12           |         | 40       | 10     |         |
| Cm                                                     | Collins silt loam<br>(occasionally<br>flooded)                                  | 39.00 | 12.2%    |                 | llw                |            |              | 110   | 800    |               | 12           |         | 40       | 10     |         |
| GrB2                                                   | Grenada silt<br>loam, 2 to 5<br>percent slopes,<br>eroded                       | 38.89 | 12.2%    |                 | lle                |            |              | 75    | 600    | 4             | 8            |         | 35       | 7.5    | 2550    |
| GrD3                                                   | Grenada silt<br>loam, 8 to 12<br>percent slopes,<br>severely<br>eroded (loring) | 26.97 | 8.4%     |                 | Vle                |            |              |       |        |               |              | 5.5     |          |        |         |
| GrD2                                                   | Grenada silt<br>loam, 8 to 12<br>percent slopes,<br>eroded (loring)             | 7.54  | 2.4%     |                 | IVe                |            |              | 75    | 600    |               |              | 7       | 25       |        |         |
| Fa                                                     | Falaya silt loam<br>(occasionally<br>flooded)                                   | 5.86  | 1.8%     |                 | llw                | 7.5        | 7            | 100   | 750    |               |              |         | 40       | 9      |         |
| FI                                                     | Falaya silt<br>loam, local<br>alluvium<br>(occasionally<br>flooded)             | 5.45  | 1.7%     |                 | llw                | 7.5        | 7            | 100   | 750    |               |              |         | 40       | 9      |         |
| Mx                                                     | Mixed alluvial<br>land (bruno<br>and Collins<br>soils, frequently<br>flooded)   | 3.46  | 1.1%     |                 | Vw                 |            |              |       |        |               |              | 4.5     |          |        |         |
| GrB3                                                   | Grenada silt<br>loam, 2 to 5<br>percent slopes,<br>severely<br>eroded           | 1.79  | 0.6%     |                 | llle               |            |              | 75    | 600    | 4             | 8            |         | 35       | 7.5    | 2550    |
| Soils data provided by USDA and NRCS. Weighted Average |                                                                                 |       |          |                 | 0.3                | 0.2        | 53.2         | 419.8 | 1.1    | 5.4           | 0.7          | 22      | 5.1      | 324.2  |         |



## Herminjard Place (320 +/- ac.) Panola Co., Miss.

## **Location and Parcel Maps**







#### Herminjard Place (320 +/- ac.) Panola Co., Miss.

### **Property Photos**



McClintock Land Company has previously entered into an agreement with the client to provide certain real estate brokerage services and acts as that client's designated agent. As such, McClintock Land is the agent of the seller. The information contained herein is from sources McClintock Land considers reliable, but is not guaranteed. This offering may be withdrawn from the market or be subject to a change in price or terms without advance notice. Seller reserves the right to reject any and all offers. Boundary lines on maps are for illustration purposes only and should not be used to define legal property lines.



Murry McClintock, Licensed Real Estate Broker

Office: 662-363-1801 murry@mcclintockland.com Cell: 662-363-6024