

For Sale

12.02 Acres | \$6,250 / AC Land Space



Harmon Rd. (12.02 Acres)

Harmon Rd. (12.02 Acres) Hopkins, South Carolina 29061

Property Highlights

- $\pm 1,480'$ of paved frontage on Harmon Road
- ±1,450' of paved frontage on Ridge Road
- ± 615 ' in width at the northern end of the property
- A couple of nice spots for homesites
- · City of Columbia Water is available
- · Less than 15 minutes from Columbia
- Sales Price: \$75,125 or \$6,250 per acre





For More Information

Tombo Milliken

O: 803 744 9852 tombo.milliken@naicolumbia.com


Tom Milliken

O: 803 744 9837 tmilliken@naicolumbia.com

Nelson Weston, III

O: 803 744 9804 nweston@naicolumbia.com

Location











ap Updated: Monday, April 26, 2021. This information submitted is not guaranteed. Although obtained from reliable sources, all information should be confirmed prior to use or reliance upon the information. This document may not be reproduced in whole or in part without the express written consent of NAI Colum














Topographical Map: USGS





FEMA Flood Zones






ap Updated: Monday, April 26, 2021. This information submitted is not guaranteed. Although obtained from reliable sources, all information should be confirmed prior to use or reliance upon the information. This document may not be reproduced in whole or in part without the express written consent of NAI Columbi









Map Unit Description (Brief, Generated)

Richland County, South Carolina

[Minor map unit components are excluded from this report]

Map unit: BaB - Blanton sand, 0 to 6 percent slopes

Component: Blanton (100%)

The Blanton component makes up 100 percent of the map unit. Slopes are 0 to 6 percent. This component is on marine terraces on sandhills. The parent material consists of sandy and loamy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is moderately well drained. Water movement in the most restrictive layer is moderately high. Available water to a depth of 60 inches is low. Shrink-swell potential is low. This soil is not flooded. It is not ponded. A seasonal zone of water saturation is at 48 inches during January, February, March, December. Organic matter content in the surface horizon is about 1 percent. Nonirrigated land capability classification is 3s. This soil does not meet hydric criteria.

Map unit: FuB - Fuquay sand, 2 to 6 percent slopes

Component: Fuquay (100%)

The Fuquay component makes up 100 percent of the map unit. Slopes are 2 to 6 percent. This component is on marine terraces on coastal plains. The parent material consists of plinthic loamy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is well drained. Water movement in the most restrictive layer is moderately low. Available water to a depth of 60 inches is low. Shrink-swell potential is low. This soil is not flooded. It is not ponded. A seasonal zone of water saturation is at 60 inches during January, February, March. Organic matter content in the surface horizon is about 1 percent. Nonirrigated land capability classification is 2s. This soil does not meet hydric criteria.

Map unit: ObB - Orangeburg loamy sand, 2 to 6 percent slopes


Component: Orangeburg (100%)

The Orangeburg component makes up 100 percent of the map unit. Slopes are 2 to 6 percent. This component is on marine terraces on coastal plains. The parent material consists of loamy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is well drained. Water movement in the most restrictive layer is moderately high. Available water to a depth of 60 inches is moderate. Shrink-swell potential is low. This soil is not flooded. It is not ponded. There is no zone of water saturation within a depth of 72 inches. Organic matter content in the surface horizon is about 1 percent. Nonirrigated land capability classification is 2e. This soil does not meet hydric criteria.

Map unit: ObC - Orangeburg loamy sand, 6 to 10 percent slopes

Component: Orangeburg (100%)

The Orangeburg component makes up 100 percent of the map unit. Slopes are 6 to 10 percent. This component is on marine terraces on coastal plains. The parent material consists of loamy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is well drained. Water movement in the most restrictive layer is moderately high. Available water to a depth of 60 inches is moderate. Shrink-swell potential is low. This soil is not flooded. It is not ponded. There is no zone of water saturation within a depth of 72 inches. Organic matter content in the surface horizon is about 1 percent. Nonirrigated land capability classification is 4e. This soil does not meet hydric criteria.



Survey Area Version: 15 Survey Area Version Date: 12/23/2013

Page 1

