

Tombo Milliken

+1 803 206 8384 tombo.milliken@naicolumbia.com

Tom Milliken

+1 803 331 6999 tmilliken@naicolumbia.com Nelson Weston

+1 803 678 7346 nweston@naicolumbia.com



TO US BY THE OWNER OF THE PROPERTY
REASON TO DOUBT ITS ACCURACY, BUT WE ERIFIED PRIOR TO PURCHASE OR LEASE.


807 Gervais Street, Suite 301
Columbia, South Carolina 29201
+1 803.254.0100
www.naicolumbia.com

Property Features

- 40± Acres for sale near Bethune, SC
- Approximately 15± Acres in 5 year-old planted Longleaf Pines
- Approximately 22± Acres in 4 year old planted Loblolly Pines
- 660± ft of paved road frontage on Mill Creek Rd
- Power along road frontage
- Property is Zoned RD-2 (Rural Resource District) in Kershaw County
- Sales Price: \$140,000 or \$3,500/ acre





































Map Unit Description (Brief, Generated)

Kershaw County Area, South Carolina

[Minor map unit components are excluded from this report]

Kershaw County Area, South Carolina

[Minor map unit components are excluded from this report]

Map unit: ApB - Alpin sand, 0 to 6 percent slopes

Component: Alpin (100%)

The Alpin component makes up 100 percent of the map unit. Slopes are 0 to 6 percent. This component is on marine terraces, coastal plains, sandhills. The parent material consists of sandy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is excessively drained. Water movement in the most restrictive layer is high. Available water to a depth of 60 inches is low. Shrink-swell potential is low. This soil is not flooded. It is not ponded. There is no zone of water saturation within a depth of 72 inches. Organic matter content in the surface horizon is about 1 percent. Nonirrigated land capability classification is 4s. This soil does not meet hydric criteria.

Map unit: BaB - Blanton sand, 0 to 6 percent slopes

Component: Blanton (100%)

The Blanton component makes up 100 percent of the map unit. Slopes are 0 to 6 percent. This component is on marine terraces, coastal plains, sandhills. The parent material consists of sandy marine deposits and loamy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is moderately well drained. Water movement in the most restrictive layer is moderately high. Available water to a depth of 60 inches is very low. Shrink-swell potential is low. This soil is not flooded. It is not ponded. A seasonal zone of water saturation is at 60 inches during January, February, March, December. Organic matter content in the surface horizon is about 1 percent. Nonirrigated land capability classification is 3s. This soil does not meet hydric criteria.

Map unit: GoA - Goldsboro loamy sand, 0 to 2 percent slopes

Component: Goldsboro (95%)

The Goldsboro component makes up 95 percent of the map unit. Slopes are 0 to 2 percent. This component is on coastal plains, marine terraces. The parent material consists of loamy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is moderately well drained. Water movement in the most restrictive layer is moderately high. Available water to a depth of 60 inches is moderate. Shrink-swell potential is low. This soil is not flooded. It is not ponded. A seasonal zone of water saturation is at 24 inches during January, February, March, April, December. Organic matter content in the surface horizon is about 1 percent. Nonirrigated land capability classification is 2w. This soil does not meet hydric criteria.

Map unit: NoA - Norfolk loamy sand, 0 to 2 percent slopes

Component: Norfolk (95%)

The Norfolk component makes up 95 percent of the map unit. Slopes are 0 to 2 percent. This component is on marine terraces, coastal plains. The parent material consists of loamy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is well drained. Water movement in the most restrictive layer is low. Available water to a depth of 60 inches is moderate. Shrink-swell potential is low. This soil is not flooded. It is not ponded. A seasonal zone of water saturation is at 48 inches during January, February, March, December. Organic matter content in the surface horizon is about 1 percent. Nonirrigated land capability classification is 1. This soil does not meet hydric criteria.

Map unit: NoB - Norfolk loamy sand, 2 to 6 percent slopes

Component: Norfolk (100%)

The Norfolk component makes up 100 percent of the map unit. Slopes are 2 to 6 percent. This component is on marine terraces, coastal plains. The parent material consists of loamy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is well drained. Water movement in the most restrictive layer is low. Available water to a depth of 60 inches is moderate. Shrink-swell potential is low. This soil is not flooded. It is not ponded. A seasonal zone of water saturation is at 48 inches during January, February, March, December. Organic matter content in the surface horizon is about 1 percent. Nonirrigated land capability classification is 2e. This soil does not meet hydric criteria.

Map unit: Ra - Rains sandy loam


Component: Rains (100%)

The Rains component makes up 100 percent of the map unit. Slopes are 0 to 2 percent. This component is on coastal plains, marine terraces. The parent material consists of loamy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is poorly drained. Water movement in the most restrictive layer is moderately high. Available water to a depth of 60 inches is low. Shrink-swell potential is low. This soil is not flooded. It is not ponded. A seasonal zone of water saturation is at 0 inches during January, February, March, April, November, December. Organic matter content in the surface horizon is about 4 percent. Nonirrigated land capability classification is 3w. This soil meets hydric criteria.

Map unit: WaB - Wagram sand, 0 to 6 percent slopes

Component: Wagram (100%)

The Wagram component makes up 100 percent of the map unit. Slopes are 2 to 6 percent. This component is on coastal plains, marine terraces. The parent material consists of loamy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is well drained. Water movement in the most restrictive layer is moderately high. Available water to a depth of 60 inches is low. Shrink-swell potential is low. This soil is not flooded. It is not ponded. There is no zone of water saturation within a depth of 72 inches. Organic matter content in the surface horizon is about 1 percent. Nonirrigated land capability classification is 2s. This soil does not meet hydric criteria.



Survey Area Version: 14 Survey Area Version Date: 12/23/2013