

FSA Tillable Soils Map





State: Iowa County: Story

8-84N-23W Location: Township: Milford

Acres: 146.27 8/18/2023 Date:







Soils data provided by USDA and NRCS.

Area Symbol: IA169, Soil Area Version: 34										
Code	Soil Description	Acres	Percent of field	CSR2 Legend	Non-Irr Class *c	*i Corn Bu	*i Soybeans Bu	CSR2**	CSR	*n NCCPI Overall
138B	Clarion loam, 2 to 6 percent slopes	39.65	27.1%		lle	225.6	65.4	89	86	83
507	Canisteo clay loam, 0 to 2 percent slopes	23.46	16.0%		llw	224.0	65.0	84	84	81
L107	Webster clay loam, Bemis moraine, 0 to 2 percent slopes	21.37	14.6%		llw	0.0	0.0	88		84
107	Webster clay loam, 0 to 2 percent slopes	14.74	10.1%		llw	224.0	65.0	86	89	82
L62C2	Storden loam, Bemis moraine, 6 to 10 percent slopes, moderately eroded	13.87	9.5%		Ille	0.0	0.0	64		66
55	Nicollet clay loam, 1 to 3 percent slopes	13.86	9.5%		lw	233.6	67.7	89	94	81
95	Harps clay loam, 0 to 2 percent slopes	9.64	6.6%		llw	198.4	57.5	72	66	82
638C2	Clarion-Storden complex, 6 to 10 percent slopes, moderately eroded	4.66	3.2%		Ille	176.0	51.0	75	61	71
6	Okoboji silty clay loam, 0 to 1 percent slopes	3.72	2.5%		IIIw	177.6	51.5	59	59	76
90	Okoboji mucky silt loam, 0 to 1 percent slopes	1.30	0.9%		IIIw	185.6	53.8	56	62	73
Weighted Average					2.07	166.6	48.3	82.8	*-	*n 80.2

^{**}IA has updated the CSR values for each county to CSR2.

Soils data provided by USDA and NRCS.

^{*-} CSR weighted average cannot be calculated on the current soils data, use prior data version for csr values.

^{*}i Yield data provided by the ISPAID Database version 8.1.1 developed by IA State University.

^{*}n: The aggregation method is "Weighted Average using all components"
*c: Using Capabilities Class Dominant Condition Aggregation Method