

Tillable Soils Map





State: **lowa** County: **Story**

Location: 8-84N-23W

Township: **Milford**Acres: **106.63**Date: **8/18/2023**







Soils data provided by USDA and NRCS.

Area Symbol: IA169, Soil Area Version: 34										
Code	Soil Description	Acres	Percent of field	CSR2 Legend	Non-Irr Class *c	*i Corn Bu	*i Soybeans Bu	CSR2**	CSR	*n NCCPI Overall
138B	Clarion loam, 2 to 6 percent slopes	35.86	33.6%		lle	225.6	65.4	89	86	83
107	Webster clay loam, 0 to 2 percent slopes	14.73	13.8%		llw	224.0	65.0	86	89	82
L107	Webster clay loam, Bemis moraine, 0 to 2 percent slopes	14.13	13.3%		llw	0.0	0.0	88		84
L62C2	Storden loam, Bemis moraine, 6 to 10 percent slopes, moderately eroded	13.66	12.8%		IIIe	0.0	0.0	64		66
507	Canisteo clay loam, 0 to 2 percent slopes	10.26	9.6%		llw	224.0	65.0	84	84	81
55	Nicollet clay loam, 1 to 3 percent slopes	9.00	8.4%		lw	233.6	67.7	89	94	81
638C2	Clarion-Storden complex, 6 to 10 percent slopes, moderately eroded	4.66	4.4%		IIIe	176.0	51.0	75	61	71
95	Harps clay loam, 0 to 2 percent slopes	3.03	2.8%		llw	198.4	57.5	72	66	82
90	Okoboji mucky silt loam, 0 to 1 percent slopes	1.30	1.2%		IIIw	185.6	53.8	56	62	73
Weighted Average						163.7	47.5	83.3	*-	*n 79.8

^{**}IA has updated the CSR values for each county to CSR2.

Soils data provided by USDA and NRCS.

^{*-} CSR weighted average cannot be calculated on the current soils data, use prior data version for csr values.

^{*}i Yield data provided by the ISPAID Database version 8.1.1 developed by IA State University.

^{*}n: The aggregation method is "Weighted Average using all components"

^{*}c: Using Capabilities Class Dominant Condition Aggregation Method