Soils Map





Area Syn	nbol: IA179, Soil Area Version: 30									
Code	Soil Description	Acres	Percent of field	CSR2 Legend	Non-Irr Class *c	*i Corn Bu	*i Soybeans Bu	CSR2**	CSR	*n NCCPI Soybeans
425D2	Keswick loam, 9 to 14 percent slopes, moderately eroded	13.99	22.9%		IVe	88.0	25.5	9	5	38
65F	Lindley loam, 18 to 25 percent slopes	13.99	22.9%		VIIe	115.2	33.4	17	10	50
730B	Nodaway-Cantril complex, 2 to 5 percent slopes	9.23	15.1%		llw	209.6	60.8	80	63	67
65E2	Lindley loam, 14 to 18 percent slopes, moderately eroded	8.83	14.5%		Vle	134.4	39.0	29	28	55
S732C2	Weller silty clay loam, 5 to 9 percent slopes, moderately eroded	7.09	11.6%		llle	0.0	0.0	59		71
269	Humeston silt loam, 0 to 2 percent slopes, occasionally flooded	3.67	6.0%		IIIw	80.0	23.2	70	58	79
716	Lawson-Quiver-Nodaway complex, 0 to 2 percent slopes, occasionally flooded	2.31	3.8%		llw	80.0	23.2	78		87
594D2	Galland loam, heavy loess, 9 to 14 percent slopes, moderately eroded	1.49	2.4%		IVe	88.0	25.5	19	5	51
594E2	Galland loam, heavy loess, 14 to 18 percent slopes, moderately eroded	0.46	0.8%		Vle	88.0	25.5	6	5	39
	Weighted Average					108.3	31.4	36.8	*-	*n 56.1

**IA has updated the CSR values for each county to CSR2.

*- CSR weighted average cannot be calculated on the current soils data, use prior data version for csr values.

*i Yield data provided by the ISPAID Database version 8.1.1 developed by IA State University.

*n: The aggregation method is "Weighted Average using all components" *c: Using Capabilities Class Dominant Condition Aggregation Method

Soils data provided by USDA and NRCS.