


BARKER RANCH NORTH

WHEELER COUNTY, TEXAS



**COLDWELL BANKER
COMMERCIAL
FIRST EQUITY**



RICHARD HEFLEY

COLDWELL BANKER FIRST EQUITY, REALTORS

A lifetime of farming and 38 years of real estate experience have given Richard Hefley an insight on how to find the perfect land for his clients.

Working with the people of Texas and Oklahoma for decades, Hefley has seen the growth and advances that they have made in the farm and ranch industry. He enjoys being a part of that growth.

"One thing that I believe about the people of Texas and Oklahoma is that they embody the pioneer spirit and look forward to new developments," says Hefley. "I like working with these pioneering individuals, because they are always striving to adapt new technologies."

Much like his clients, Hefley also strives to adapt to new technologies. Working with his associate, Hefley researches every property he lists for sale, and compiles a book containing soil and topographic maps along with production possibilities for each property. These books are available to prospective buyers to help them determine the best land for what they want to accomplish in the areas of farming, hunting and ranching.

In addition to working with farmers and ranchers, Hefley has devoted his time to being a member of national associations to better his skills and knowledge for those he serves.

Hefley attends the Realtors Land Institute (RLI) National Land Conference every year to inform clients about national real estate trends that affect our area. He is an Accredited Land Consultant (ALC). ALC designations are held by real estate professionals that are the highest performing and most experienced land specialists; there are only 500 in the entire nation.

RICHARD HEFLEY, ALC, LICENSED IN TEXAS AND OKLAHOMA, ASSOCIATE BROKER OF COLDWELL BANKER FIRST EQUITY, REALTORS

5701 Time Square Blvd., Suite 190, Amarillo, TX 79119
806.898.1786 • richardhefley@sbcglobal.net




COUNTY RD S

COUNTY RD S

COUNTY RD 18



COUNTY RD S

BARKER RANCH NORTH



WHEELER COUNTY TEXAS

GENERAL ROADS MAP




In general, address numbers INCREASE going to the North and East at approximately one mile per one hundred block. AS address numbers INCREASE, EVEN addresses will be on the RIGHT, and ODD addresses will be on the LEFT.

This map is produced to aid emergency service providers in locating 9-1-1 calls. The information on the map, having been obtained from a variety of sources, is believed to be accurate. Please report corrections, omissions and changes to the Panhandle Regional Planning Commission's 9-1-1 network at www.panhandlemaps.com



**PANHANDLE
REGIONAL
PLANNING
COMMISSION**

March 2011



Accurate Map of Adams Blvd. Drive



United States
Department of
Agriculture


NRCS

Natural
Resources
Conservation
Service

A product of the National
Cooperative Soil Survey,
a joint effort of the United
States Department of
Agriculture and other
Federal agencies, State
agencies including the
Agricultural Experiment
Stations, and local
participants

Custom Soil Resource Report for **Wheeler County, Texas**

BARKER RANCH NORTH



April 23, 2024

Preface

Soil surveys contain information that affects land use planning in survey areas. They highlight soil limitations that affect various land uses and provide information about the properties of the soils in the survey areas. Soil surveys are designed for many different users, including farmers, ranchers, foresters, agronomists, urban planners, community officials, engineers, developers, builders, and home buyers. Also, conservationists, teachers, students, and specialists in recreation, waste disposal, and pollution control can use the surveys to help them understand, protect, or enhance the environment.

Various land use regulations of Federal, State, and local governments may impose special restrictions on land use or land treatment. Soil surveys identify soil properties that are used in making various land use or land treatment decisions. The information is intended to help the land users identify and reduce the effects of soil limitations on various land uses. The landowner or user is responsible for identifying and complying with existing laws and regulations.

Although soil survey information can be used for general farm, local, and wider area planning, onsite investigation is needed to supplement this information in some cases. Examples include soil quality assessments (<http://www.nrcs.usda.gov/wps/portal/nrcs/main/soils/health/>) and certain conservation and engineering applications. For more detailed information, contact your local USDA Service Center (<https://offices.sc.egov.usda.gov/locator/app?agency=nrcs>) or your NRCS State Soil Scientist (http://www.nrcs.usda.gov/wps/portal/nrcs/detail/soils/contactus/?cid=nrcs142p2_053951).

Great differences in soil properties can occur within short distances. Some soils are seasonally wet or subject to flooding. Some are too unstable to be used as a foundation for buildings or roads. Clayey or wet soils are poorly suited to use as septic tank absorption fields. A high water table makes a soil poorly suited to basements or underground installations.

The National Cooperative Soil Survey is a joint effort of the United States Department of Agriculture and other Federal agencies, State agencies including the Agricultural Experiment Stations, and local agencies. The Natural Resources Conservation Service (NRCS) has leadership for the Federal part of the National Cooperative Soil Survey.

Information about soils is updated periodically. Updated information is available through the NRCS Web Soil Survey, the site for official soil survey information.

The U.S. Department of Agriculture (USDA) prohibits discrimination in all its programs and activities on the basis of race, color, national origin, age, disability, and where applicable, sex, marital status, familial status, parental status, religion, sexual orientation, genetic information, political beliefs, reprisal, or because all or a part of an individual's income is derived from any public assistance program. (Not all prohibited bases apply to all programs.) Persons with disabilities who require

alternative means for communication of program information (Braille, large print, audiotape, etc.) should contact USDA's TARGET Center at (202) 720-2600 (voice and TDD). To file a complaint of discrimination, write to USDA, Director, Office of Civil Rights, 1400 Independence Avenue, S.W., Washington, D.C. 20250-9410 or call (800) 795-3272 (voice) or (202) 720-6382 (TDD). USDA is an equal opportunity provider and employer.

Contents

Preface	2
How Soil Surveys Are Made	5
Soil Map	8
Soil Map.....	9
Legend.....	10
Map Unit Legend.....	11
Map Unit Descriptions.....	11
Wheeler County, Texas.....	14
CaB—Carey silt loam, 1 to 3 percent slopes.....	14
CoB—Cobb loamy fine sand, loamy substratum 1 to 3 percent slopes.....	15
CoC—Cobb loamy fine sand, loamy substratum, 3 to 5 percent slopes.....	16
DIB—Devol loamy sand, 0 to 3 percent slopes.....	17
DID—Devol loamy sand, 3 to 8 percent slopes.....	19
GfB—Grandfield fine sandy loam, 1 to 3 percent slopes.....	21
Gu—Guadalupe fine sandy loam, 0 to 2 percent slopes, occasionally flooded.....	22
LuC—Lutie silt loam, 3 to 5 percent slopes.....	24
OuD—Obaro-Quinlan association, 5 to 12 percent slopes.....	25
PaC—Paducah loam, 3 to 5 percent slopes.....	27
Sw—Sweetwater silty clay loam, saline, 0 to 2 percent slopes, occasionally flooded.....	29
Soil Information for All Uses	31
Suitabilities and Limitations for Use.....	31
Vegetative Productivity.....	31
Range Production (Normal Year) (BARKER RANCH NORTH).....	31
Yields of Non-Irrigated Crops (Component): Cotton lint (Lbs) (BARKER RANCH NORTH).....	35
Yields of Non-Irrigated Crops (Component): Grain sorghum (Bu) (BARKER RANCH NORTH).....	39
Yields of Non-Irrigated Crops (Component): Wheat (Bu) (BARKER RANCH NORTH).....	43
References	47

How Soil Surveys Are Made

Soil surveys are made to provide information about the soils and miscellaneous areas in a specific area. They include a description of the soils and miscellaneous areas and their location on the landscape and tables that show soil properties and limitations affecting various uses. Soil scientists observed the steepness, length, and shape of the slopes; the general pattern of drainage; the kinds of crops and native plants; and the kinds of bedrock. They observed and described many soil profiles. A soil profile is the sequence of natural layers, or horizons, in a soil. The profile extends from the surface down into the unconsolidated material in which the soil formed or from the surface down to bedrock. The unconsolidated material is devoid of roots and other living organisms and has not been changed by other biological activity.

Currently, soils are mapped according to the boundaries of major land resource areas (MLRAs). MLRAs are geographically associated land resource units that share common characteristics related to physiography, geology, climate, water resources, soils, biological resources, and land uses (USDA, 2006). Soil survey areas typically consist of parts of one or more MLRA.

The soils and miscellaneous areas in a survey area occur in an orderly pattern that is related to the geology, landforms, relief, climate, and natural vegetation of the area. Each kind of soil and miscellaneous area is associated with a particular kind of landform or with a segment of the landform. By observing the soils and miscellaneous areas in the survey area and relating their position to specific segments of the landform, a soil scientist develops a concept, or model, of how they were formed. Thus, during mapping, this model enables the soil scientist to predict with a considerable degree of accuracy the kind of soil or miscellaneous area at a specific location on the landscape.

Commonly, individual soils on the landscape merge into one another as their characteristics gradually change. To construct an accurate soil map, however, soil scientists must determine the boundaries between the soils. They can observe only a limited number of soil profiles. Nevertheless, these observations, supplemented by an understanding of the soil-vegetation-landscape relationship, are sufficient to verify predictions of the kinds of soil in an area and to determine the boundaries.

Soil scientists recorded the characteristics of the soil profiles that they studied. They noted soil color, texture, size and shape of soil aggregates, kind and amount of rock fragments, distribution of plant roots, reaction, and other features that enable them to identify soils. After describing the soils in the survey area and determining their properties, the soil scientists assigned the soils to taxonomic classes (units). Taxonomic classes are concepts. Each taxonomic class has a set of soil characteristics with precisely defined limits. The classes are used as a basis for comparison to classify soils systematically. Soil taxonomy, the system of taxonomic classification used in the United States, is based mainly on the kind and character of soil properties and the arrangement of horizons within the profile. After the soil

scientists classified and named the soils in the survey area, they compared the individual soils with similar soils in the same taxonomic class in other areas so that they could confirm data and assemble additional data based on experience and research.

The objective of soil mapping is not to delineate pure map unit components; the objective is to separate the landscape into landforms or landform segments that have similar use and management requirements. Each map unit is defined by a unique combination of soil components and/or miscellaneous areas in predictable proportions. Some components may be highly contrasting to the other components of the map unit. The presence of minor components in a map unit in no way diminishes the usefulness or accuracy of the data. The delineation of such landforms and landform segments on the map provides sufficient information for the development of resource plans. If intensive use of small areas is planned, onsite investigation is needed to define and locate the soils and miscellaneous areas.

Soil scientists make many field observations in the process of producing a soil map. The frequency of observation is dependent upon several factors, including scale of mapping, intensity of mapping, design of map units, complexity of the landscape, and experience of the soil scientist. Observations are made to test and refine the soil-landscape model and predictions and to verify the classification of the soils at specific locations. Once the soil-landscape model is refined, a significantly smaller number of measurements of individual soil properties are made and recorded. These measurements may include field measurements, such as those for color, depth to bedrock, and texture, and laboratory measurements, such as those for content of sand, silt, clay, salt, and other components. Properties of each soil typically vary from one point to another across the landscape.

Observations for map unit components are aggregated to develop ranges of characteristics for the components. The aggregated values are presented. Direct measurements do not exist for every property presented for every map unit component. Values for some properties are estimated from combinations of other properties.

While a soil survey is in progress, samples of some of the soils in the area generally are collected for laboratory analyses and for engineering tests. Soil scientists interpret the data from these analyses and tests as well as the field-observed characteristics and the soil properties to determine the expected behavior of the soils under different uses. Interpretations for all of the soils are field tested through observation of the soils in different uses and under different levels of management. Some interpretations are modified to fit local conditions, and some new interpretations are developed to meet local needs. Data are assembled from other sources, such as research information, production records, and field experience of specialists. For example, data on crop yields under defined levels of management are assembled from farm records and from field or plot experiments on the same kinds of soil.

Predictions about soil behavior are based not only on soil properties but also on such variables as climate and biological activity. Soil conditions are predictable over long periods of time, but they are not predictable from year to year. For example, soil scientists can predict with a fairly high degree of accuracy that a given soil will have a high water table within certain depths in most years, but they cannot predict that a high water table will always be at a specific level in the soil on a specific date.

After soil scientists located and identified the significant natural bodies of soil in the survey area, they drew the boundaries of these bodies on aerial photographs and


Custom Soil Resource Report

identified each as a specific map unit. Aerial photographs show trees, buildings, fields, roads, and rivers, all of which help in locating boundaries accurately.

Soil Map

The soil map section includes the soil map for the defined area of interest, a list of soil map units on the map and extent of each map unit, and cartographic symbols displayed on the map. Also presented are various metadata about data used to produce the map, and a description of each soil map unit.

Custom Soil Resource Report Soil Map



Custom Soil Resource Report

MAP LEGEND

Area of Interest (AOI)

 Area of Interest (AOI)

Soils

 Soil Map Unit Polygons

 Soil Map Unit Lines

 Soil Map Unit Points

Special Point Features

 Blowout

 Borrow Pit

 Clay Spot

 Closed Depression

 Gravel Pit

 Gravelly Spot

 Landfill

 Lava Flow

 Marsh or swamp

 Mine or Quarry

 Miscellaneous Water

 Perennial Water

 Rock Outcrop

 Saline Spot

 Sandy Spot

 Severely Eroded Spot

 Sinkhole

 Slide or Slip

 Sodic Spot

 Spoil Area

 Stony Spot

 Very Stony Spot

 Wet Spot

 Other

 Special Line Features

Water Features

 Streams and Canals

Transportation

 Rails

 Interstate Highways

 US Routes

 Major Roads

 Local Roads

Background

 Aerial Photography

MAP INFORMATION

The soil surveys that comprise your AOI were mapped at 1:24,000.

Warning: Soil Map may not be valid at this scale.

Enlargement of maps beyond the scale of mapping can cause misunderstanding of the detail of mapping and accuracy of soil line placement. The maps do not show the small areas of contrasting soils that could have been shown at a more detailed scale.

Please rely on the bar scale on each map sheet for map measurements.

Source of Map: Natural Resources Conservation Service
Web Soil Survey URL:
Coordinate System: Web Mercator (EPSG:3857)

Maps from the Web Soil Survey are based on the Web Mercator projection, which preserves direction and shape but distorts distance and area. A projection that preserves area, such as the Albers equal-area conic projection, should be used if more accurate calculations of distance or area are required.

This product is generated from the USDA-NRCS certified data as of the version date(s) listed below.

Soil Survey Area: Wheeler County, Texas
Survey Area Data: Version 19, Sep 5, 2023

Soil map units are labeled (as space allows) for map scales 1:50,000 or larger.

Date(s) aerial images were photographed: Nov 23, 2021—Dec 5, 2021

The orthophoto or other base map on which the soil lines were compiled and digitized probably differs from the background imagery displayed on these maps. As a result, some minor shifting of map unit boundaries may be evident.

Map Unit Legend

Map Unit Symbol	Map Unit Name	Acres in AOI	Percent of AOI
CaB	Carey silt loam, 1 to 3 percent slopes	5.9	1.8%
CoB	Cobb loamy fine sand, loamy substratum 1 to 3 percent slopes	55.4	17.4%
CoC	Cobb loamy fine sand, loamy substratum, 3 to 5 percent slopes	22.3	7.0%
DIB	Devol loamy sand, 0 to 3 percent slopes	44.0	13.8%
DID	Devol loamy sand, 3 to 8 percent slopes	82.5	25.8%
GfB	Grandfield fine sandy loam, 1 to 3 percent slopes	12.6	3.9%
Gu	Guadalupe fine sandy loam, 0 to 2 percent slopes, occasionally flooded	0.7	0.2%
LuC	Lutie silt loam, 3 to 5 percent slopes	19.8	6.2%
OuD	Obaro-Quinlan association, 5 to 12 percent slopes	70.2	22.0%
PaC	Paducah loam, 3 to 5 percent slopes	5.0	1.6%
Sw	Sweetwater silty clay loam, saline, 0 to 2 percent slopes, occasionally flooded	0.8	0.3%
Totals for Area of Interest		319.2	100.0%

Map Unit Descriptions

The map units delineated on the detailed soil maps in a soil survey represent the soils or miscellaneous areas in the survey area. The map unit descriptions, along with the maps, can be used to determine the composition and properties of a unit.

A map unit delineation on a soil map represents an area dominated by one or more major kinds of soil or miscellaneous areas. A map unit is identified and named according to the taxonomic classification of the dominant soils. Within a taxonomic class there are precisely defined limits for the properties of the soils. On the landscape, however, the soils are natural phenomena, and they have the characteristic variability of all natural phenomena. Thus, the range of some observed properties may extend beyond the limits defined for a taxonomic class. Areas of soils of a single taxonomic class rarely, if ever, can be mapped without including areas of other taxonomic classes. Consequently, every map unit is made

up of the soils or miscellaneous areas for which it is named and some minor components that belong to taxonomic classes other than those of the major soils.

Most minor soils have properties similar to those of the dominant soil or soils in the map unit, and thus they do not affect use and management. These are called noncontrasting, or similar, components. They may or may not be mentioned in a particular map unit description. Other minor components, however, have properties and behavioral characteristics divergent enough to affect use or to require different management. These are called contrasting, or dissimilar, components. They generally are in small areas and could not be mapped separately because of the scale used. Some small areas of strongly contrasting soils or miscellaneous areas are identified by a special symbol on the maps. If included in the database for a given area, the contrasting minor components are identified in the map unit descriptions along with some characteristics of each. A few areas of minor components may not have been observed, and consequently they are not mentioned in the descriptions, especially where the pattern was so complex that it was impractical to make enough observations to identify all the soils and miscellaneous areas on the landscape.

The presence of minor components in a map unit in no way diminishes the usefulness or accuracy of the data. The objective of mapping is not to delineate pure taxonomic classes but rather to separate the landscape into landforms or landform segments that have similar use and management requirements. The delineation of such segments on the map provides sufficient information for the development of resource plans. If intensive use of small areas is planned, however, onsite investigation is needed to define and locate the soils and miscellaneous areas.

An identifying symbol precedes the map unit name in the map unit descriptions. Each description includes general facts about the unit and gives important soil properties and qualities.

Soils that have profiles that are almost alike make up a *soil series*. Except for differences in texture of the surface layer, all the soils of a series have major horizons that are similar in composition, thickness, and arrangement.

Soils of one series can differ in texture of the surface layer, slope, stoniness, salinity, degree of erosion, and other characteristics that affect their use. On the basis of such differences, a soil series is divided into *soil phases*. Most of the areas shown on the detailed soil maps are phases of soil series. The name of a soil phase commonly indicates a feature that affects use or management. For example, Alpha silt loam, 0 to 2 percent slopes, is a phase of the Alpha series.

Some map units are made up of two or more major soils or miscellaneous areas. These map units are complexes, associations, or undifferentiated groups.

A *complex* consists of two or more soils or miscellaneous areas in such an intricate pattern or in such small areas that they cannot be shown separately on the maps. The pattern and proportion of the soils or miscellaneous areas are somewhat similar in all areas. Alpha-Beta complex, 0 to 6 percent slopes, is an example.

An *association* is made up of two or more geographically associated soils or miscellaneous areas that are shown as one unit on the maps. Because of present or anticipated uses of the map units in the survey area, it was not considered practical or necessary to map the soils or miscellaneous areas separately. The pattern and relative proportion of the soils or miscellaneous areas are somewhat similar. Alpha-Beta association, 0 to 2 percent slopes, is an example.

Custom Soil Resource Report

An *undifferentiated group* is made up of two or more soils or miscellaneous areas that could be mapped individually but are mapped as one unit because similar interpretations can be made for use and management. The pattern and proportion of the soils or miscellaneous areas in a mapped area are not uniform. An area can be made up of only one of the major soils or miscellaneous areas, or it can be made up of all of them. Alpha and Beta soils, 0 to 2 percent slopes, is an example.

Some surveys include *miscellaneous areas*. Such areas have little or no soil material and support little or no vegetation. Rock outcrop is an example.

Wheeler County, Texas

CaB—Carey silt loam, 1 to 3 percent slopes

Map Unit Setting

National map unit symbol: 2t6pd
Elevation: 1,000 to 2,500 feet
Mean annual precipitation: 20 to 28 inches
Mean annual air temperature: 57 to 64 degrees F
Frost-free period: 180 to 230 days
Farmland classification: All areas are prime farmland

Map Unit Composition

Carey and similar soils: 85 percent
Minor components: 15 percent
Estimates are based on observations, descriptions, and transects of the mapunit.

Description of Carey

Setting

Landform: Hillslopes on hills
Landform position (two-dimensional): Backslope
Landform position (three-dimensional): Side slope
Down-slope shape: Convex
Across-slope shape: Convex
Parent material: Sandy and silty residuum weathered from sandstone and siltstone

Typical profile

Ap - 0 to 14 inches: silt loam
Bt - 14 to 36 inches: silty clay loam
Bk - 36 to 50 inches: loam
Cd - 50 to 80 inches: very fine sandy loam

Properties and qualities

Slope: 1 to 3 percent
Depth to restrictive feature: 36 to 56 inches to densic bedrock
Drainage class: Well drained
Runoff class: Low
Capacity of the most limiting layer to transmit water (Ksat): Very low to moderately low (0.00 to 0.06 in/hr)
Depth to water table: More than 80 inches
Frequency of flooding: None
Frequency of ponding: None
Calcium carbonate, maximum content: 20 percent
Gypsum, maximum content: 2 percent
Maximum salinity: Nonsaline (0.0 to 1.0 mmhos/cm)
Sodium adsorption ratio, maximum: 4.0
Available water supply, 0 to 60 inches: Moderate (about 7.3 inches)

Interpretive groups

Land capability classification (irrigated): None specified
Land capability classification (nonirrigated): 2e
Hydrologic Soil Group: B
Ecological site: R078CY056OK - Loamy Upland
Hydric soil rating: No

Minor Components

Woodward

Percent of map unit: 5 percent
Landform: Hills
Landform position (two-dimensional): Backslope
Landform position (three-dimensional): Side slope
Down-slope shape: Convex
Across-slope shape: Convex
Ecological site: R078CY056OK - Loamy Upland
Hydric soil rating: No

Quinlan

Percent of map unit: 5 percent
Landform: Hills
Landform position (two-dimensional): Backslope
Landform position (three-dimensional): Side slope
Down-slope shape: Convex
Across-slope shape: Convex
Ecological site: R078CY083OK - Shallow Upland
Hydric soil rating: No

St. paul

Percent of map unit: 5 percent
Landform: Paleoterraces
Landform position (three-dimensional): Tread
Down-slope shape: Convex
Across-slope shape: Linear
Ecological site: R078CY056OK - Loamy Upland
Hydric soil rating: No

CoB—Cobb loamy fine sand, loamy substratum 1 to 3 percent slopes

Map Unit Setting

National map unit symbol: djfh
Elevation: 1,000 to 2,200 feet
Mean annual precipitation: 20 to 28 inches
Mean annual air temperature: 61 to 68 degrees F
Frost-free period: 210 to 240 days
Farmland classification: Not prime farmland

Map Unit Composition

Cobb and similar soils: 100 percent
Estimates are based on observations, descriptions, and transects of the mapunit.

Description of Cobb

Setting

Landform: Plains
Down-slope shape: Convex

Custom Soil Resource Report

Across-slope shape: Linear

Parent material: Loamy residuum weathered from sandstone

Typical profile

H1 - 0 to 16 inches: loamy fine sand

H2 - 16 to 38 inches: sandy clay loam

H3 - 38 to 65 inches: bedrock

Properties and qualities

Slope: 1 to 3 percent

Depth to restrictive feature: 20 to 40 inches to paralithic bedrock

Drainage class: Well drained

Runoff class: Medium

Capacity of the most limiting layer to transmit water (Ksat): Moderately low to moderately high (0.06 to 0.20 in/hr)

Depth to water table: More than 80 inches

Frequency of flooding: None

Frequency of ponding: None

Calcium carbonate, maximum content: 5 percent

Maximum salinity: Nonsaline to very slightly saline (0.0 to 2.0 mmhos/cm)

Available water supply, 0 to 60 inches: Low (about 4.1 inches)

Interpretive groups

Land capability classification (irrigated): 3e

Land capability classification (nonirrigated): 3e

Hydrologic Soil Group: C

Ecological site: R078CY110TX - Sandy Loam 23-31" PZ

Hydric soil rating: No

CoC—Cobb loamy fine sand, loamy substratum, 3 to 5 percent slopes

Map Unit Setting

National map unit symbol: djfj

Elevation: 1,000 to 2,200 feet

Mean annual precipitation: 20 to 28 inches

Mean annual air temperature: 61 to 68 degrees F

Frost-free period: 210 to 240 days

Farmland classification: Not prime farmland

Map Unit Composition

Cobb and similar soils: 100 percent

Estimates are based on observations, descriptions, and transects of the mapunit.

Description of Cobb

Setting

Landform: Plains

Down-slope shape: Convex

Across-slope shape: Linear

Parent material: Loamy residuum weathered from sandstone

Typical profile

H1 - 0 to 9 inches: loamy fine sand

Custom Soil Resource Report

H2 - 9 to 38 inches: sandy clay loam

H3 - 38 to 65 inches: bedrock

Properties and qualities

Slope: 3 to 5 percent

Depth to restrictive feature: 20 to 40 inches to paralithic bedrock

Drainage class: Well drained

Runoff class: Medium

Capacity of the most limiting layer to transmit water (Ksat): Moderately low to moderately high (0.06 to 0.20 in/hr)

Depth to water table: More than 80 inches

Frequency of flooding: None

Frequency of ponding: None

Calcium carbonate, maximum content: 5 percent

Maximum salinity: Nonsaline to very slightly saline (0.0 to 2.0 mmhos/cm)

Available water supply, 0 to 60 inches: Low (about 4.3 inches)

Interpretive groups

Land capability classification (irrigated): 4e

Land capability classification (nonirrigated): 4e

Hydrologic Soil Group: C

Ecological site: R078CY110TX - Sandy Loam 23-31" PZ

Hydric soil rating: No

DIB—Devol loamy sand, 0 to 3 percent slopes

Map Unit Setting

National map unit symbol: 2sdbh

Elevation: 1,000 to 2,500 feet

Mean annual precipitation: 22 to 31 inches

Mean annual air temperature: 57 to 64 degrees F

Frost-free period: 185 to 230 days

Farmland classification: Not prime farmland

Map Unit Composition

Devol and similar soils: 90 percent

Minor components: 10 percent

Estimates are based on observations, descriptions, and transects of the mapunit.

Description of Devol

Setting

Landform: Dunes

Landform position (two-dimensional): Backslope

Landform position (three-dimensional): Side slope, tread

Down-slope shape: Convex

Across-slope shape: Convex

Parent material: Sandy eolian deposits over coarse-loamy alluvium

Typical profile

Ap - 0 to 14 inches: loamy sand

Bt - 14 to 34 inches: fine sandy loam

Custom Soil Resource Report

BC - 34 to 46 inches: fine sandy loam

C - 46 to 80 inches: loamy fine sand

Properties and qualities

Slope: 0 to 3 percent

Depth to restrictive feature: More than 80 inches

Drainage class: Well drained

Runoff class: Very low

Capacity of the most limiting layer to transmit water (Ksat): High (1.98 to 5.95 in/hr)

Depth to water table: More than 80 inches

Frequency of flooding: None

Frequency of ponding: None

Calcium carbonate, maximum content: 1 percent

Maximum salinity: Nonsaline (0.0 to 0.1 mmhos/cm)

Available water supply, 0 to 60 inches: Moderate (about 6.2 inches)

Interpretive groups

Land capability classification (irrigated): 3e

Land capability classification (nonirrigated): 3e

Hydrologic Soil Group: A

Ecological site: R078CY014OK - Rolling Sands

Hydric soil rating: No

Minor Components

Grandfield

Percent of map unit: 5 percent

Landform: Sand sheets on paleoterraces

Landform position (three-dimensional): Tread

Down-slope shape: Convex

Across-slope shape: Convex

Ecological site: R078CY110TX - Sandy Loam 23-31" PZ

Hydric soil rating: No

Eda

Percent of map unit: 3 percent

Landform: Dunes

Landform position (two-dimensional): Backslope

Landform position (three-dimensional): Side slope

Down-slope shape: Convex

Across-slope shape: Convex

Ecological site: R078CY014OK - Rolling Sands

Hydric soil rating: No

Grandmore

Percent of map unit: 1 percent

Landform: Sand sheets on paleoterraces

Landform position (three-dimensional): Tread

Down-slope shape: Convex

Across-slope shape: Convex

Ecological site: R078CY014OK - Rolling Sands

Hydric soil rating: No

Carwile

Percent of map unit: 1 percent

Landform: Depressions

Custom Soil Resource Report

Down-slope shape: Concave
Across-slope shape: Concave
Ecological site: R078CY025OK - Depressional Upland
Hydric soil rating: Yes

DID—Devol loamy sand, 3 to 8 percent slopes

Map Unit Setting

National map unit symbol: 2sdbl
Elevation: 980 to 2,590 feet
Mean annual precipitation: 22 to 31 inches
Mean annual air temperature: 57 to 64 degrees F
Frost-free period: 185 to 230 days
Farmland classification: Not prime farmland

Map Unit Composition

Devol and similar soils: 80 percent
Minor components: 20 percent
Estimates are based on observations, descriptions, and transects of the mapunit.

Description of Devol

Setting

Landform: Dunes
Landform position (two-dimensional): Backslope
Landform position (three-dimensional): Side slope, tread
Down-slope shape: Convex
Across-slope shape: Convex
Parent material: Coarse-loamy alluvium over sandy eolian deposits

Typical profile

A - 0 to 10 inches: loamy sand
Bt - 10 to 31 inches: fine sandy loam
C - 31 to 80 inches: loamy fine sand

Properties and qualities

Slope: 3 to 8 percent
Depth to restrictive feature: More than 80 inches
Drainage class: Well drained
Runoff class: Low
Capacity of the most limiting layer to transmit water (Ksat): High (2.00 to 6.00 in/hr)
Depth to water table: More than 80 inches
Frequency of flooding: None
Frequency of ponding: None
Available water supply, 0 to 60 inches: Moderate (about 6.4 inches)

Interpretive groups

Land capability classification (irrigated): None specified
Land capability classification (nonirrigated): 4e
Hydrologic Soil Group: A

Custom Soil Resource Report

Ecological site: R078CY014OK - Rolling Sands

Hydric soil rating: No

Minor Components

Grandfield

Percent of map unit: 8 percent

Landform: Sand sheets on paleoterraces

Landform position (three-dimensional): Riser

Down-slope shape: Convex

Across-slope shape: Convex

Ecological site: R078CY110TX - Sandy Loam 23-31" PZ

Hydric soil rating: No

Eda

Percent of map unit: 4 percent

Landform: Dunes

Landform position (two-dimensional): Backslope

Landform position (three-dimensional): Side slope

Down-slope shape: Convex

Across-slope shape: Convex

Ecological site: R078CY014OK - Rolling Sands

Hydric soil rating: No

Dill

Percent of map unit: 3 percent

Landform: Hillslopes on hills

Landform position (two-dimensional): Backslope

Down-slope shape: Convex

Across-slope shape: Convex

Ecological site: R078CY110TX - Sandy Loam 23-31" PZ

Hydric soil rating: No

Grandmore

Percent of map unit: 2 percent

Landform: Sand sheets on paleoterraces

Landform position (three-dimensional): Tread

Down-slope shape: Convex

Across-slope shape: Convex

Ecological site: R078CY014OK - Rolling Sands

Hydric soil rating: No

Hardeman

Percent of map unit: 2 percent

Landform: Stream terraces

Landform position (three-dimensional): Tread

Down-slope shape: Convex

Across-slope shape: Convex

Ecological site: R078CY110TX - Sandy Loam 23-31" PZ

Hydric soil rating: No

Carwile

Percent of map unit: 1 percent

Landform: Depressions

Down-slope shape: Concave

Across-slope shape: Concave

Ecological site: R078CY025OK - Depressional Upland

Hydric soil rating: Yes

GfB—Grandfield fine sandy loam, 1 to 3 percent slopes

Map Unit Setting

National map unit symbol: 2txy9

Elevation: 1,000 to 2,200 feet

Mean annual precipitation: 21 to 28 inches

Mean annual air temperature: 57 to 64 degrees F

Frost-free period: 180 to 230 days

Farmland classification: Farmland of statewide importance, if irrigated

Map Unit Composition

Grandfield and similar soils: 90 percent

Minor components: 10 percent

Estimates are based on observations, descriptions, and transects of the mapunit.

Description of Grandfield

Setting

Landform: Sand sheets on paleoterraces

Landform position (three-dimensional): Tread

Down-slope shape: Convex

Across-slope shape: Linear

Parent material: Loamy alluvium and/or eolian deposits

Typical profile

Ap - 0 to 9 inches: fine sandy loam

Bt1 - 9 to 23 inches: fine sandy loam

Bt2 - 23 to 57 inches: sandy clay loam

BC - 57 to 80 inches: fine sandy loam

Properties and qualities

Slope: 1 to 3 percent

Depth to restrictive feature: More than 80 inches

Drainage class: Well drained

Runoff class: Low

Capacity of the most limiting layer to transmit water (Ksat): Moderately high to high
(0.57 to 1.98 in/hr)

Depth to water table: More than 80 inches

Frequency of flooding: None

Frequency of ponding: None

Calcium carbonate, maximum content: 5 percent

Available water supply, 0 to 60 inches: Moderate (about 8.1 inches)

Interpretive groups

Land capability classification (irrigated): 2e

Land capability classification (nonirrigated): 2e

Hydrologic Soil Group: B

Ecological site: R078CY110TX - Sandy Loam 23-31" PZ

Hydric soil rating: No

Minor Components

Devol

Percent of map unit: 4 percent
Landform: Dunes on sand sheets on stream terraces
Landform position (two-dimensional): Backslope
Landform position (three-dimensional): Side slope
Down-slope shape: Convex
Across-slope shape: Convex
Ecological site: R078CY110TX - Sandy Loam 23-31" PZ
Hydric soil rating: No

Delwin

Percent of map unit: 3 percent
Landform: Sand sheets
Landform position (three-dimensional): Tread
Down-slope shape: Convex
Across-slope shape: Convex
Ecological site: R078CY017OK - Deep Sand Shrubland
Hydric soil rating: No

Grandmore

Percent of map unit: 2 percent
Landform: Sand sheets on paleoterraces
Landform position (three-dimensional): Tread
Down-slope shape: Concave
Across-slope shape: Concave
Ecological site: R078CY110TX - Sandy Loam 23-31" PZ
Hydric soil rating: No

Carwile

Percent of map unit: 1 percent
Landform: Depressions on sand sheets
Landform position (two-dimensional): Backslope, toeslope, shoulder
Landform position (three-dimensional): Side slope, tread
Down-slope shape: Convex, concave
Across-slope shape: Convex, concave
Ecological site: R078CY025OK - Depressional Upland
Hydric soil rating: Yes

Gu—Guadalupe fine sandy loam, 0 to 2 percent slopes, occasionally flooded

Map Unit Setting

National map unit symbol: 2w83x
Elevation: 1,900 to 4,600 feet
Mean annual precipitation: 15 to 26 inches
Mean annual air temperature: 56 to 61 degrees F
Frost-free period: 180 to 200 days
Farmland classification: Farmland of statewide importance, if irrigated

Map Unit Composition

Guadalupe, occasionally flooded, and similar soils: 82 percent

Minor components: 18 percent

Estimates are based on observations, descriptions, and transects of the mapunit.

Description of Guadalupe, Occasionally Flooded

Setting

Landform: Flood plains

Landform position (three-dimensional): Tread

Down-slope shape: Linear

Across-slope shape: Linear

Parent material: Holocene calcareous sandy and loamy alluvium

Typical profile

A - 0 to 12 inches: fine sandy loam

Bw - 12 to 25 inches: fine sandy loam

Bk - 25 to 38 inches: fine sandy loam

Ck - 38 to 80 inches: loamy sand

Properties and qualities

Slope: 0 to 2 percent

Depth to restrictive feature: More than 80 inches

Drainage class: Well drained

Runoff class: Negligible

*Capacity of the most limiting layer to transmit water (Ksat): Moderately high to high
(0.57 to 5.95 in/hr)*

Depth to water table: More than 80 inches

Frequency of flooding: Occasional

Frequency of ponding: None

Calcium carbonate, maximum content: 4 percent

Maximum salinity: Nonsaline (0.1 to 0.5 mmhos/cm)

Available water supply, 0 to 60 inches: Moderate (about 6.7 inches)

Interpretive groups

Land capability classification (irrigated): 3e

Land capability classification (nonirrigated): 3c

Hydrologic Soil Group: A

Ecological site: R077EY065TX - Sandy Bottomland 16-24" PZ

Hydric soil rating: No

Minor Components

Touzalin, occasionally flooded

Percent of map unit: 8 percent

Landform: Flood plains

Landform position (three-dimensional): Tread

Down-slope shape: Linear

Across-slope shape: Linear

Ecological site: R077EY065TX - Sandy Bottomland 16-24" PZ

Hydric soil rating: No

Wespur, occasionally flooded

Percent of map unit: 5 percent

Landform: Flood plains

Landform position (three-dimensional): Tread

Custom Soil Resource Report

Down-slope shape: Linear
Across-slope shape: Linear
Ecological site: R077EY058TX - Loamy Bottomland 16-24" PZ
Hydric soil rating: No

Sweetwater, occasionally flooded

Percent of map unit: 3 percent
Landform: Flood plains
Landform position (three-dimensional): Tread
Down-slope shape: Linear
Across-slope shape: Concave
Ecological site: R077EY571TX - Wet Bottomland 16-24" PZ
Hydric soil rating: Yes

Dreyfoos

Percent of map unit: 2 percent
Landform: Dunes on flood plains
Landform position (three-dimensional): Tread
Down-slope shape: Convex
Across-slope shape: Convex
Ecological site: R077EY063TX - Sand Hills 16-24" PZ
Hydric soil rating: No

LuC—Lutie silt loam, 3 to 5 percent slopes

Map Unit Setting

National map unit symbol: djg8
Elevation: 2,000 to 2,600 feet
Mean annual precipitation: 22 to 26 inches
Mean annual air temperature: 59 to 63 degrees F
Frost-free period: 200 to 220 days
Farmland classification: All areas are prime farmland

Map Unit Composition

Lutie and similar soils: 100 percent
Estimates are based on observations, descriptions, and transects of the mapunit.

Description of Lutie

Setting

Landform: Plains
Down-slope shape: Convex
Across-slope shape: Linear
Parent material: Loamy alluvium

Typical profile

H1 - 0 to 11 inches: silt loam
H2 - 11 to 30 inches: silty clay loam
H3 - 30 to 80 inches: silty clay loam

Properties and qualities

Slope: 3 to 5 percent
Depth to restrictive feature: More than 80 inches
Drainage class: Well drained
Runoff class: Low
Capacity of the most limiting layer to transmit water (Ksat): Moderately high to high
(0.57 to 1.98 in/hr)
Depth to water table: More than 80 inches
Frequency of flooding: None
Frequency of ponding: None
Calcium carbonate, maximum content: 40 percent
Available water supply, 0 to 60 inches: High (about 10.1 inches)

Interpretive groups

Land capability classification (irrigated): None specified
Land capability classification (nonirrigated): 4e
Hydrologic Soil Group: B
Ecological site: R078BY081TX - Loamy Upland 19-26" PZ
Hydric soil rating: No

OuD—Obaro-Quinlan association, 5 to 12 percent slopes

Map Unit Setting

National map unit symbol: 2t6q3
Elevation: 1,000 to 2,500 feet
Mean annual precipitation: 20 to 28 inches
Mean annual air temperature: 57 to 64 degrees F
Frost-free period: 180 to 230 days
Farmland classification: Not prime farmland

Map Unit Composition

Obaro and similar soils: 55 percent
Quinlan and similar soils: 30 percent
Minor components: 15 percent
Estimates are based on observations, descriptions, and transects of the mapunit.

Description of Obaro

Setting

Landform: Knolls, ridges
Landform position (two-dimensional): Summit, shoulder
Landform position (three-dimensional): Crest, side slope
Down-slope shape: Convex, linear
Across-slope shape: Convex
Parent material: Loamy residuum weathered from calcareous sandstone

Typical profile

A - 0 to 7 inches: loam
Bw - 7 to 17 inches: loam
Bk - 17 to 34 inches: loam
Cr - 34 to 60 inches: bedrock

Properties and qualities

Slope: 5 to 12 percent
Depth to restrictive feature: 29 to 38 inches to paralithic bedrock
Drainage class: Well drained
Runoff class: Medium
Capacity of the most limiting layer to transmit water (Ksat): Moderately high (0.20 to 0.57 in/hr)
Depth to water table: More than 80 inches
Frequency of flooding: None
Frequency of ponding: None
Calcium carbonate, maximum content: 25 percent
Maximum salinity: Nonsaline to very slightly saline (0.0 to 2.0 mmhos/cm)
Available water supply, 0 to 60 inches: Low (about 5.0 inches)

Interpretive groups

Land capability classification (irrigated): None specified
Land capability classification (nonirrigated): 6e
Hydrologic Soil Group: C
Ecological site: R078BY081TX - Loamy Upland 19-26" PZ
Hydric soil rating: No

Description of Quinlan

Setting

Landform: Hillslopes on hills
Landform position (two-dimensional): Shoulder, backslope
Landform position (three-dimensional): Side slope
Down-slope shape: Convex
Across-slope shape: Convex, linear
Parent material: Loamy residuum weathered from calcareous sandstone

Typical profile

A - 0 to 7 inches: loam
Bw - 7 to 12 inches: loam
Cd - 12 to 60 inches: very fine sandy loam

Properties and qualities

Slope: 5 to 12 percent
Depth to restrictive feature: 8 to 18 inches to densic bedrock
Drainage class: Well drained
Runoff class: High
Capacity of the most limiting layer to transmit water (Ksat): Moderately high (0.20 to 0.57 in/hr)
Depth to water table: More than 80 inches
Frequency of flooding: None
Frequency of ponding: None
Calcium carbonate, maximum content: 15 percent
Gypsum, maximum content: 2 percent
Maximum salinity: Nonsaline (0.0 to 1.0 mmhos/cm)
Sodium adsorption ratio, maximum: 4.0
Available water supply, 0 to 60 inches: Very low (about 2.1 inches)

Interpretive groups

Land capability classification (irrigated): None specified
Land capability classification (nonirrigated): 6e
Hydrologic Soil Group: D

Custom Soil Resource Report

Ecological site: R078BY089TX - Shallow 19-26" PZ

Hydric soil rating: No

Minor Components

Aspermont

Percent of map unit: 5 percent

Landform: Hillslopes

Landform position (two-dimensional): Shoulder, backslope

Landform position (three-dimensional): Side slope

Down-slope shape: Convex

Across-slope shape: Convex

Ecological site: R078BY079TX - Loamy 19-26" PZ

Hydric soil rating: No

Carey

Percent of map unit: 5 percent

Landform: Hillslopes on hills

Landform position (two-dimensional): Backslope

Landform position (three-dimensional): Side slope

Down-slope shape: Convex

Across-slope shape: Convex

Ecological site: R078BY081TX - Loamy Upland 19-26" PZ

Hydric soil rating: No

Grandfield

Percent of map unit: 5 percent

Landform: Sand sheets on paleoterraces

Landform position (three-dimensional): Tread

Down-slope shape: Convex

Across-slope shape: Linear

Ecological site: R078CY110TX - Sandy Loam 23-31" PZ

Hydric soil rating: No

PaC—Paducah loam, 3 to 5 percent slopes

Map Unit Setting

National map unit symbol: 2wrnd

Elevation: 1,600 to 2,500 feet

Mean annual precipitation: 21 to 25 inches

Mean annual air temperature: 59 to 64 degrees F

Frost-free period: 200 to 225 days

Farmland classification: All areas are prime farmland

Map Unit Composition

Paducah and similar soils: 90 percent

Minor components: 10 percent

Estimates are based on observations, descriptions, and transects of the mapunit.

Description of Paducah

Setting

Landform: Terraces, hillslopes

Landform position (two-dimensional): Shoulder, backslope

Landform position (three-dimensional): Side slope, riser

Down-slope shape: Convex

Across-slope shape: Convex

Parent material: Loamy alluvium and/or residuum weathered from calcareous sandstone and/or calcareous siltstone

Typical profile

Ap - 0 to 8 inches: loam

Bt - 8 to 27 inches: clay loam

Bk - 27 to 72 inches: very fine sandy loam

Cd - 72 to 90 inches: very fine sandy loam

Properties and qualities

Slope: 3 to 5 percent

Depth to restrictive feature: 60 to 80 inches to densic bedrock

Drainage class: Well drained

Runoff class: Low

Capacity of the most limiting layer to transmit water (Ksat): Moderately high to high (0.20 to 1.98 in/hr)

Depth to water table: More than 80 inches

Frequency of flooding: None

Frequency of ponding: None

Calcium carbonate, maximum content: 20 percent

Gypsum, maximum content: 2 percent

Maximum salinity: Nonsaline to very slightly saline (0.0 to 2.0 mmhos/cm)

Sodium adsorption ratio, maximum: 3.0

Available water supply, 0 to 60 inches: High (about 9.2 inches)

Interpretive groups

Land capability classification (irrigated): 3e

Land capability classification (nonirrigated): 3e

Hydrologic Soil Group: B

Ecological site: R078BY081TX - Loamy Upland 19-26" PZ

Hydric soil rating: No

Minor Components

Woodward

Percent of map unit: 5 percent

Landform: Hillslopes

Landform position (two-dimensional): Shoulder, backslope

Landform position (three-dimensional): Side slope

Down-slope shape: Convex

Across-slope shape: Convex

Ecological site: R078BY081TX - Loamy Upland 19-26" PZ

Hydric soil rating: No

Obaro

Percent of map unit: 4 percent

Landform: Knolls on hillslopes

Landform position (two-dimensional): Backslope, shoulder

Custom Soil Resource Report

Landform position (three-dimensional): Side slope
Down-slope shape: Convex
Across-slope shape: Convex
Ecological site: R078BY081TX - Loamy Upland 19-26" PZ
Hydric soil rating: No

Quinlan

Percent of map unit: 1 percent
Landform: Knolls on hillslopes
Landform position (two-dimensional): Backslope, shoulder
Landform position (three-dimensional): Side slope
Down-slope shape: Convex
Across-slope shape: Convex
Ecological site: R078BY089TX - Shallow 19-26" PZ
Hydric soil rating: No

Sw—Sweetwater silty clay loam, saline, 0 to 2 percent slopes, occasionally flooded

Map Unit Setting

National map unit symbol: 23d06
Elevation: 1,800 to 3,200 feet
Mean annual precipitation: 16 to 26 inches
Mean annual air temperature: 56 to 59 degrees F
Frost-free period: 180 to 210 days
Farmland classification: Not prime farmland

Map Unit Composition

Sweetwater, saline, occasionally flooded, and similar soils: 90 percent
Minor components: 10 percent
Estimates are based on observations, descriptions, and transects of the mapunit.

Description of Sweetwater, Saline, Occasionally Flooded

Setting

Landform: Sloughs, flood plains
Landform position (three-dimensional): Tread
Down-slope shape: Concave
Across-slope shape: Concave
Parent material: Holocene fine-loamy alluvium over calcareous sandy alluvium

Typical profile

A1 - 0 to 14 inches: silty clay loam
A2 - 14 to 24 inches: clay loam
Cn - 24 to 80 inches: sand

Properties and qualities

Slope: 0 to 2 percent
Depth to restrictive feature: 15 to 34 inches to strongly contrasting textural stratification
Drainage class: Poorly drained

Custom Soil Resource Report

Runoff class: Negligible

Capacity of the most limiting layer to transmit water (Ksat): Moderately high to high
(0.57 to 1.98 in/hr)

Depth to water table: About 6 to 20 inches

Frequency of flooding: Occasional

Frequency of ponding: None

Calcium carbonate, maximum content: 8 percent

Maximum salinity: Nonsaline to moderately saline (0.5 to 8.0 mmhos/cm)

Sodium adsorption ratio, maximum: 30.0

Available water supply, 0 to 60 inches: Low (about 4.2 inches)

Interpretive groups

Land capability classification (irrigated): 4w

Land capability classification (nonirrigated): 4w

Hydrologic Soil Group: B/D

Ecological site: R077EY571TX - Wet Bottomland 16-24" PZ

Hydric soil rating: Yes

Minor Components

Persimmon, occasionally flooded

Percent of map unit: 7 percent

Landform: Flood plains

Landform position (three-dimensional): Tread

Down-slope shape: Linear

Across-slope shape: Linear

Ecological site: R077EY571TX - Wet Bottomland 16-24" PZ

Hydric soil rating: No

Touzalin, occasionally flooded

Percent of map unit: 3 percent

Landform: Flood plains

Landform position (three-dimensional): Tread

Down-slope shape: Linear

Across-slope shape: Linear

Ecological site: R077EY065TX - Sandy Bottomland 16-24" PZ

Hydric soil rating: No

Soil Information for All Uses

Suitabilities and Limitations for Use

The Suitabilities and Limitations for Use section includes various soil interpretations displayed as thematic maps with a summary table for the soil map units in the selected area of interest. A single value or rating for each map unit is generated by aggregating the interpretive ratings of individual map unit components. This aggregation process is defined for each interpretation.

Vegetative Productivity


Vegetative productivity includes estimates of potential vegetative production for a variety of land uses, including cropland, forestland, hayland, pastureland, horticulture and rangeland. In the underlying database, some states maintain crop yield data by individual map unit component. Other states maintain the data at the map unit level. Attributes are included for both, although only one or the other is likely to contain data for any given geographic area. For other land uses, productivity data is shown only at the map unit component level. Examples include potential crop yields under irrigated and nonirrigated conditions, forest productivity, forest site index, and total rangeland production under of normal, favorable and unfavorable conditions.

Range Production (Normal Year) (BARKER RANCH NORTH)

Total range production is the amount of vegetation that can be expected to grow annually in a well managed area that is supporting the potential natural plant community. It includes all vegetation, whether or not it is palatable to grazing animals. It includes the current year's growth of leaves, twigs, and fruits of woody plants. It does not include the increase in stem diameter of trees and shrubs. It is expressed in pounds per acre of air-dry vegetation. In a normal year, growing conditions are about average. Yields are adjusted to a common percent of air-dry moisture content.

In areas that have similar climate and topography, differences in the kind and amount of vegetation produced on rangeland are closely related to the kind of soil. Effective management is based on the relationship between the soils and vegetation and water.

Custom Soil Resource Report
Map—Range Production (Normal Year) (BARKER RANCH NORTH)



MAP LEGEND

Area of Interest (AOI)

 Area of Interest (AOI)

Soils

Soil Rating Polygons

 <= 1804
 > 1804 and <= 2100
 > 2100 and <= 2517
 > 2517 and <= 2930
 > 2930 and <= 4080
 Not rated or not available

Soil Rating Lines

 <= 1804
 > 1804 and <= 2100
 > 2100 and <= 2517
 > 2517 and <= 2930
 > 2930 and <= 4080
 Not rated or not available

Soil Rating Points

 <= 1804
 > 1804 and <= 2100
 > 2100 and <= 2517
 > 2517 and <= 2930
 > 2930 and <= 4080
 Not rated or not available

Water Features

 Streams and Canals

Transportation

 Rails
 Interstate Highways
 US Routes
 Major Roads
 Local Roads

Background

 Aerial Photography

MAP INFORMATION

The soil surveys that comprise your AOI were mapped at 1:24,000.

Warning: Soil Map may not be valid at this scale.

Enlargement of maps beyond the scale of mapping can cause misunderstanding of the detail of mapping and accuracy of soil line placement. The maps do not show the small areas of contrasting soils that could have been shown at a more detailed scale.

Please rely on the bar scale on each map sheet for map measurements.

Source of Map: Natural Resources Conservation Service
 Web Soil Survey URL:
 Coordinate System: Web Mercator (EPSG:3857)

Maps from the Web Soil Survey are based on the Web Mercator projection, which preserves direction and shape but distorts distance and area. A projection that preserves area, such as the Albers equal-area conic projection, should be used if more accurate calculations of distance or area are required.

This product is generated from the USDA-NRCS certified data as of the version date(s) listed below.

Soil Survey Area: Wheeler County, Texas
 Survey Area Data: Version 19, Sep 5, 2023

Soil map units are labeled (as space allows) for map scales 1:50,000 or larger.

Date(s) aerial images were photographed: Nov 23, 2021—Dec 5, 2021

The orthophoto or other base map on which the soil lines were compiled and digitized probably differs from the background imagery displayed on these maps. As a result, some minor shifting of map unit boundaries may be evident.

Table—Range Production (Normal Year) (BARKER RANCH NORTH)

Map unit symbol	Map unit name	Rating (pounds per acre per year)	Acres in AOI	Percent of AOI
CaB	Carey silt loam, 1 to 3 percent slopes	4080	5.9	1.8%
CoB	Cobb loamy fine sand, loamy substratum 1 to 3 percent slopes	2500	55.4	17.4%
CoC	Cobb loamy fine sand, loamy substratum, 3 to 5 percent slopes	2500	22.3	7.0%
DIB	Devol loamy sand, 0 to 3 percent slopes	2462	44.0	13.8%
DID	Devol loamy sand, 3 to 8 percent slopes	2517	82.5	25.8%
GfB	Grandfield fine sandy loam, 1 to 3 percent slopes	2762	12.6	3.9%
Gu	Guadalupe fine sandy loam, 0 to 2 percent slopes, occasionally flooded	1804	0.7	0.2%
LuC	Lutie silt loam, 3 to 5 percent slopes	1800	19.8	6.2%
OuD	Obaro-Quinlan association, 5 to 12 percent slopes	2100	70.2	22.0%
PaC	Paducah loam, 3 to 5 percent slopes	1800	5.0	1.6%
Sw	Sweetwater silty clay loam, saline, 0 to 2 percent slopes, occasionally flooded	2930	0.8	0.3%
Totals for Area of Interest			319.2	100.0%

Rating Options—Range Production (Normal Year) (BARKER RANCH NORTH)*Units of Measure:* pounds per acre per year*Aggregation Method:* Weighted Average*Component Percent Cutoff:* None Specified*Tie-break Rule:* Higher*Interpret Nulls as Zero:* Yes

Yields of Non-Irrigated Crops (Component): Cotton lint (Lbs) (BARKER RANCH NORTH)

These are the estimated average yields per acre that can be expected of selected nonirrigated crops under a high level of management. In any given year, yields may be higher or lower than those indicated because of variations in rainfall and other climatic factors.

In the database, some states maintain crop yield data by individual map unit component and others maintain the data at the map unit level. Attributes are included in this application for both, although only one or the other is likely to contain data for any given geographic area. This attribute uses data maintained at the map unit component level.


The yields are actually recorded as three separate values in the database. A low value and a high value indicate the range for the soil component. A "representative" value indicates the expected value for the component. For these yields, only the representative value is used.

The yields are based mainly on the experience and records of farmers, conservationists, and extension agents. Available yield data from nearby areas and results of field trials and demonstrations also are considered.

The management needed to obtain the indicated yields of the various crops depends on the kind of soil and the crop. Management can include drainage, erosion control, and protection from flooding; the proper planting and seeding rates; suitable high-yielding crop varieties; appropriate and timely tillage; control of weeds, plant diseases, and harmful insects; favorable soil reaction and optimum levels of nitrogen, phosphorus, potassium, and trace elements for each crop; effective use of crop residue, barnyard manure, and green manure crops; and harvesting that ensures the smallest possible loss.

The estimated yields reflect the productive capacity of each soil for the selected crop. Yields are likely to increase as new production technology is developed. The productivity of a given soil compared with that of other soils, however, is not likely to change.

Custom Soil Resource Report
 Map—Yields of Non-Irrigated Crops (Component): Cotton lint (Lbs) (BARKER RANCH NORTH)



MAP LEGEND

Area of Interest (AOI)

 Area of Interest (AOI)

Soils

Soil Rating Polygons

 ≤ 150.00
 > 150.00 and ≤ 180.00
 > 180.00 and ≤ 200.00
 > 200.00 and ≤ 244.75
 > 244.75 and ≤ 286.75
 Not rated or not available

Soil Rating Lines

 ≤ 150.00
 > 150.00 and ≤ 180.00
 > 180.00 and ≤ 200.00
 > 200.00 and ≤ 244.75
 > 244.75 and ≤ 286.75
 Not rated or not available

Soil Rating Points

 ≤ 150.00
 > 150.00 and ≤ 180.00
 > 180.00 and ≤ 200.00
 > 200.00 and ≤ 244.75
 > 244.75 and ≤ 286.75
 Not rated or not available

Water Features

 Streams and Canals

Transportation

 Rails
 Interstate Highways
 US Routes
 Major Roads
 Local Roads

Background

 Aerial Photography

MAP INFORMATION

The soil surveys that comprise your AOI were mapped at 1:24,000.

Warning: Soil Map may not be valid at this scale.

Enlargement of maps beyond the scale of mapping can cause misunderstanding of the detail of mapping and accuracy of soil line placement. The maps do not show the small areas of contrasting soils that could have been shown at a more detailed scale.

Please rely on the bar scale on each map sheet for map measurements.

Source of Map: Natural Resources Conservation Service
 Web Soil Survey URL:
 Coordinate System: Web Mercator (EPSG:3857)

Maps from the Web Soil Survey are based on the Web Mercator projection, which preserves direction and shape but distorts distance and area. A projection that preserves area, such as the Albers equal-area conic projection, should be used if more accurate calculations of distance or area are required.

This product is generated from the USDA-NRCS certified data as of the version date(s) listed below.

Soil Survey Area: Wheeler County, Texas
 Survey Area Data: Version 19, Sep 5, 2023

Soil map units are labeled (as space allows) for map scales 1:50,000 or larger.

Date(s) aerial images were photographed: Nov 23, 2021—Dec 5, 2021

The orthophoto or other base map on which the soil lines were compiled and digitized probably differs from the background imagery displayed on these maps. As a result, some minor shifting of map unit boundaries may be evident.

Table—Yields of Non-Irrigated Crops (Component): Cotton lint (Lbs) (BARKER RANCH NORTH)

Map unit symbol	Map unit name	Rating	Acres in AOI	Percent of AOI
CaB	Carey silt loam, 1 to 3 percent slopes	233.75	5.9	1.8%
CoB	Cobb loamy fine sand, loamy substratum 1 to 3 percent slopes	200.00	55.4	17.4%
CoC	Cobb loamy fine sand, loamy substratum, 3 to 5 percent slopes		22.3	7.0%
DIB	Devol loamy sand, 0 to 3 percent slopes	244.75	44.0	13.8%
DID	Devol loamy sand, 3 to 8 percent slopes	177.00	82.5	25.8%
GfB	Grandfield fine sandy loam, 1 to 3 percent slopes	286.75	12.6	3.9%
Gu	Guadalupe fine sandy loam, 0 to 2 percent slopes, occasionally flooded		0.7	0.2%
LuC	Lutie silt loam, 3 to 5 percent slopes	150.00	19.8	6.2%
OuD	Obaro-Quinlan association, 5 to 12 percent slopes	143.75	70.2	22.0%
PaC	Paducah loam, 3 to 5 percent slopes	180.00	5.0	1.6%
Sw	Sweetwater silty clay loam, saline, 0 to 2 percent slopes, occasionally flooded		0.8	0.3%
Totals for Area of Interest			319.2	100.0%

Rating Options—Yields of Non-Irrigated Crops (Component): Cotton lint (Lbs) (BARKER RANCH NORTH)*Crop:* Cotton lint*Yield Units:* Lbs*Aggregation Method:* Weighted Average*Component Percent Cutoff:* None Specified*Tie-break Rule:* Higher*Interpret Nulls as Zero:* Yes

Yields of Non-Irrigated Crops (Component): Grain sorghum (Bu) (BARKER RANCH NORTH)

These are the estimated average yields per acre that can be expected of selected nonirrigated crops under a high level of management. In any given year, yields may be higher or lower than those indicated because of variations in rainfall and other climatic factors.

In the database, some states maintain crop yield data by individual map unit component and others maintain the data at the map unit level. Attributes are included in this application for both, although only one or the other is likely to contain data for any given geographic area. This attribute uses data maintained at the map unit component level.


The yields are actually recorded as three separate values in the database. A low value and a high value indicate the range for the soil component. A "representative" value indicates the expected value for the component. For these yields, only the representative value is used.

The yields are based mainly on the experience and records of farmers, conservationists, and extension agents. Available yield data from nearby areas and results of field trials and demonstrations also are considered.

The management needed to obtain the indicated yields of the various crops depends on the kind of soil and the crop. Management can include drainage, erosion control, and protection from flooding; the proper planting and seeding rates; suitable high-yielding crop varieties; appropriate and timely tillage; control of weeds, plant diseases, and harmful insects; favorable soil reaction and optimum levels of nitrogen, phosphorus, potassium, and trace elements for each crop; effective use of crop residue, barnyard manure, and green manure crops; and harvesting that ensures the smallest possible loss.

The estimated yields reflect the productive capacity of each soil for the selected crop. Yields are likely to increase as new production technology is developed. The productivity of a given soil compared with that of other soils, however, is not likely to change.

Custom Soil Resource Report
 Map—Yields of Non-Irrigated Crops (Component): Grain sorghum (Bu) (BARKER RANCH NORTH)



MAP LEGEND

Area of Interest (AOI)

 Area of Interest (AOI)

Soils

Soil Rating Polygons

 ≤ 13.12
 > 13.12 and ≤ 24.80
 > 24.80 and ≤ 25.20
 > 25.20 and ≤ 29.55
 > 29.55 and ≤ 29.75
 Not rated or not available

Soil Rating Lines

 ≤ 13.12
 > 13.12 and ≤ 24.80
 > 24.80 and ≤ 25.20
 > 25.20 and ≤ 29.55
 > 29.55 and ≤ 29.75
 Not rated or not available

Soil Rating Points

 ≤ 13.12
 > 13.12 and ≤ 24.80
 > 24.80 and ≤ 25.20
 > 25.20 and ≤ 29.55
 > 29.55 and ≤ 29.75
 Not rated or not available

Water Features

 Streams and Canals

Transportation

 Rails
 Interstate Highways
 US Routes
 Major Roads
 Local Roads

Background

 Aerial Photography

MAP INFORMATION

The soil surveys that comprise your AOI were mapped at 1:24,000.

Warning: Soil Map may not be valid at this scale.

Enlargement of maps beyond the scale of mapping can cause misunderstanding of the detail of mapping and accuracy of soil line placement. The maps do not show the small areas of contrasting soils that could have been shown at a more detailed scale.

Please rely on the bar scale on each map sheet for map measurements.

Source of Map: Natural Resources Conservation Service
 Web Soil Survey URL:
 Coordinate System: Web Mercator (EPSG:3857)

Maps from the Web Soil Survey are based on the Web Mercator projection, which preserves direction and shape but distorts distance and area. A projection that preserves area, such as the Albers equal-area conic projection, should be used if more accurate calculations of distance or area are required.

This product is generated from the USDA-NRCS certified data as of the version date(s) listed below.

Soil Survey Area: Wheeler County, Texas
 Survey Area Data: Version 19, Sep 5, 2023

Soil map units are labeled (as space allows) for map scales 1:50,000 or larger.

Date(s) aerial images were photographed: Nov 23, 2021—Dec 5, 2021

The orthophoto or other base map on which the soil lines were compiled and digitized probably differs from the background imagery displayed on these maps. As a result, some minor shifting of map unit boundaries may be evident.

Table—Yields of Non-Irrigated Crops (Component): Grain sorghum (Bu) (BARKER RANCH NORTH)

Map unit symbol	Map unit name	Rating	Acres in AOI	Percent of AOI
CaB	Carey silt loam, 1 to 3 percent slopes	29.75	5.9	1.8%
CoB	Cobb loamy fine sand, loamy substratum 1 to 3 percent slopes	25.00	55.4	17.4%
CoC	Cobb loamy fine sand, loamy substratum, 3 to 5 percent slopes		22.3	7.0%
DIB	Devol loamy sand, 0 to 3 percent slopes	29.45	44.0	13.8%
DID	Devol loamy sand, 3 to 8 percent slopes	25.10	82.5	25.8%
GfB	Grandfield fine sandy loam, 1 to 3 percent slopes	29.55	12.6	3.9%
Gu	Guadalupe fine sandy loam, 0 to 2 percent slopes, occasionally flooded	13.12	0.7	0.2%
LuC	Lutie silt loam, 3 to 5 percent slopes	25.00	19.8	6.2%
OuD	Obaro-Quinlan association, 5 to 12 percent slopes	24.80	70.2	22.0%
PaC	Paducah loam, 3 to 5 percent slopes	25.20	5.0	1.6%
Sw	Sweetwater silty clay loam, saline, 0 to 2 percent slopes, occasionally flooded		0.8	0.3%
Totals for Area of Interest			319.2	100.0%

Rating Options—Yields of Non-Irrigated Crops (Component): Grain sorghum (Bu) (BARKER RANCH NORTH)*Crop:* Grain sorghum*Yield Units:* Bu*Aggregation Method:* Weighted Average*Component Percent Cutoff:* None Specified*Tie-break Rule:* Higher*Interpret Nulls as Zero:* Yes

Yields of Non-Irrigated Crops (Component): Wheat (Bu) (BARKER RANCH NORTH)

These are the estimated average yields per acre that can be expected of selected nonirrigated crops under a high level of management. In any given year, yields may be higher or lower than those indicated because of variations in rainfall and other climatic factors.

In the database, some states maintain crop yield data by individual map unit component and others maintain the data at the map unit level. Attributes are included in this application for both, although only one or the other is likely to contain data for any given geographic area. This attribute uses data maintained at the map unit component level.


The yields are actually recorded as three separate values in the database. A low value and a high value indicate the range for the soil component. A "representative" value indicates the expected value for the component. For these yields, only the representative value is used.

The yields are based mainly on the experience and records of farmers, conservationists, and extension agents. Available yield data from nearby areas and results of field trials and demonstrations also are considered.

The management needed to obtain the indicated yields of the various crops depends on the kind of soil and the crop. Management can include drainage, erosion control, and protection from flooding; the proper planting and seeding rates; suitable high-yielding crop varieties; appropriate and timely tillage; control of weeds, plant diseases, and harmful insects; favorable soil reaction and optimum levels of nitrogen, phosphorus, potassium, and trace elements for each crop; effective use of crop residue, barnyard manure, and green manure crops; and harvesting that ensures the smallest possible loss.

The estimated yields reflect the productive capacity of each soil for the selected crop. Yields are likely to increase as new production technology is developed. The productivity of a given soil compared with that of other soils, however, is not likely to change.

Custom Soil Resource Report
 Map—Yields of Non-Irrigated Crops (Component): Wheat (Bu) (BARKER RANCH NORTH)



MAP LEGEND

Area of Interest (AOI)

 Area of Interest (AOI)

Soils

Soil Rating Polygons

 ≤ 15.00
 > 15.00 and ≤ 16.30
 > 16.30 and ≤ 19.74
 > 19.74 and ≤ 24.20
 > 24.20 and ≤ 27.18
 Not rated or not available

Soil Rating Lines

 ≤ 15.00
 > 15.00 and ≤ 16.30
 > 16.30 and ≤ 19.74
 > 19.74 and ≤ 24.20
 > 24.20 and ≤ 27.18
 Not rated or not available

Soil Rating Points

 ≤ 15.00
 > 15.00 and ≤ 16.30
 > 16.30 and ≤ 19.74
 > 19.74 and ≤ 24.20
 > 24.20 and ≤ 27.18
 Not rated or not available

Water Features

 Streams and Canals

Transportation

 Rails
 Interstate Highways
 US Routes
 Major Roads
 Local Roads

Background

 Aerial Photography

MAP INFORMATION

The soil surveys that comprise your AOI were mapped at 1:24,000.

Warning: Soil Map may not be valid at this scale.

Enlargement of maps beyond the scale of mapping can cause misunderstanding of the detail of mapping and accuracy of soil line placement. The maps do not show the small areas of contrasting soils that could have been shown at a more detailed scale.

Please rely on the bar scale on each map sheet for map measurements.

Source of Map: Natural Resources Conservation Service
 Web Soil Survey URL:
 Coordinate System: Web Mercator (EPSG:3857)

Maps from the Web Soil Survey are based on the Web Mercator projection, which preserves direction and shape but distorts distance and area. A projection that preserves area, such as the Albers equal-area conic projection, should be used if more accurate calculations of distance or area are required.

This product is generated from the USDA-NRCS certified data as of the version date(s) listed below.

Soil Survey Area: Wheeler County, Texas
 Survey Area Data: Version 19, Sep 5, 2023

Soil map units are labeled (as space allows) for map scales 1:50,000 or larger.

Date(s) aerial images were photographed: Nov 23, 2021—Dec 5, 2021

The orthophoto or other base map on which the soil lines were compiled and digitized probably differs from the background imagery displayed on these maps. As a result, some minor shifting of map unit boundaries may be evident.

**Table—Yields of Non-Irrigated Crops (Component): Wheat (Bu)
(BARKER RANCH NORTH)**

Map unit symbol	Map unit name	Rating	Acres in AOI	Percent of AOI
CaB	Carey silt loam, 1 to 3 percent slopes	24.20	5.9	1.8%
CoB	Cobb loamy fine sand, loamy substratum 1 to 3 percent slopes	15.00	55.4	17.4%
CoC	Cobb loamy fine sand, loamy substratum, 3 to 5 percent slopes		22.3	7.0%
DIB	Devol loamy sand, 0 to 3 percent slopes	19.74	44.0	13.8%
DID	Devol loamy sand, 3 to 8 percent slopes	15.73	82.5	25.8%
GfB	Grandfield fine sandy loam, 1 to 3 percent slopes	27.18	12.6	3.9%
Gu	Guadalupe fine sandy loam, 0 to 2 percent slopes, occasionally flooded		0.7	0.2%
LuC	Lutie silt loam, 3 to 5 percent slopes	15.00	19.8	6.2%
OuD	Obaro-Quinlan association, 5 to 12 percent slopes	16.30	70.2	22.0%
PaC	Paducah loam, 3 to 5 percent slopes	14.40	5.0	1.6%
Sw	Sweetwater silty clay loam, saline, 0 to 2 percent slopes, occasionally flooded		0.8	0.3%
Totals for Area of Interest			319.2	100.0%

**Rating Options—Yields of Non-Irrigated Crops (Component):
Wheat (Bu) (BARKER RANCH NORTH)***Crop:* Wheat*Yield Units:* Bu*Aggregation Method:* Weighted Average*Component Percent Cutoff:* None Specified*Tie-break Rule:* Higher*Interpret Nulls as Zero:* Yes

References

- American Association of State Highway and Transportation Officials (AASHTO). 2004. Standard specifications for transportation materials and methods of sampling and testing. 24th edition.
- American Society for Testing and Materials (ASTM). 2005. Standard classification of soils for engineering purposes. ASTM Standard D2487-00.
- Cowardin, L.M., V. Carter, F.C. Golet, and E.T. LaRoe. 1979. Classification of wetlands and deep-water habitats of the United States. U.S. Fish and Wildlife Service FWS/OBS-79/31.
- Federal Register. July 13, 1994. Changes in hydric soils of the United States.
- Federal Register. September 18, 2002. Hydric soils of the United States.
- Hurt, G.W., and L.M. Vasilas, editors. Version 6.0, 2006. Field indicators of hydric soils in the United States.
- National Research Council. 1995. Wetlands: Characteristics and boundaries.
- Soil Survey Division Staff. 1993. Soil survey manual. Soil Conservation Service. U.S. Department of Agriculture Handbook 18. http://www.nrcs.usda.gov/wps/portal/nrcs/detail/national/soils/?cid=nrcs142p2_054262
- Soil Survey Staff. 1999. Soil taxonomy: A basic system of soil classification for making and interpreting soil surveys. 2nd edition. Natural Resources Conservation Service, U.S. Department of Agriculture Handbook 436. http://www.nrcs.usda.gov/wps/portal/nrcs/detail/national/soils/?cid=nrcs142p2_053577
- Soil Survey Staff. 2010. Keys to soil taxonomy. 11th edition. U.S. Department of Agriculture, Natural Resources Conservation Service. http://www.nrcs.usda.gov/wps/portal/nrcs/detail/national/soils/?cid=nrcs142p2_053580
- Tiner, R.W., Jr. 1985. Wetlands of Delaware. U.S. Fish and Wildlife Service and Delaware Department of Natural Resources and Environmental Control, Wetlands Section.
- United States Army Corps of Engineers, Environmental Laboratory. 1987. Corps of Engineers wetlands delineation manual. Waterways Experiment Station Technical Report Y-87-1.
- United States Department of Agriculture, Natural Resources Conservation Service. National forestry manual. http://www.nrcs.usda.gov/wps/portal/nrcs/detail/soils/home/?cid=nrcs142p2_053374
- United States Department of Agriculture, Natural Resources Conservation Service. National range and pasture handbook. <http://www.nrcs.usda.gov/wps/portal/nrcs/detail/national/landuse/rangepasture/?cid=stelprdb1043084>

Custom Soil Resource Report

United States Department of Agriculture, Natural Resources Conservation Service. National soil survey handbook, title 430-VI. http://www.nrcs.usda.gov/wps/portal/nrcs/detail/soils/scientists/?cid=nrcs142p2_054242

United States Department of Agriculture, Natural Resources Conservation Service. 2006. Land resource regions and major land resource areas of the United States, the Caribbean, and the Pacific Basin. U.S. Department of Agriculture Handbook 296. http://www.nrcs.usda.gov/wps/portal/nrcs/detail/national/soils/?cid=nrcs142p2_053624

United States Department of Agriculture, Soil Conservation Service. 1961. Land capability classification. U.S. Department of Agriculture Handbook 210. http://www.nrcs.usda.gov/Internet/FSE_DOCUMENTS/nrcs142p2_052290.pdf