



TILLABLE SOILS





Field borders obtained from Farm Service Agency as of 2008

FarmWorth, LLC makes no representations or warranties, express or implied, as to the accuracy of any information, data, numerical values, boundaries, or any other information generated through the use of FarmWorth.com. User is solely responsible for independently investigating and determining all information provided through FarmWorth.com prior to use and waives any and all claims against FarmWorth, LLC for any inaccuracies or inconsistencies in the information and/or data.

Code	Description	Acres	% of field	CPI	Soil Class (Non-Irr)
F1152B 📃	Balaton-Souris-Svea loams, 0 to 4 percent slopes	54.72	39.86 %	75	2
F122A	Svea-Cresbard loams, 0 to 3 percent slopes	24.77	18.04 %	76	2
F1102A 🗌	Hamerly-Wyard-Tonka complex, 0 to 3 percent slopes	15.44	11.25 %	68	2
F135A 📃	Hamerly-Cresbard loams, 0 to 3 percent slopes	9.35	6.81 %	71	2
F135B 🗌	Balaton-Cresbard loams, 3 to 6 percent slopes	6.96	5.07 %	67	2
F1012A 📕	Parnell-Vallers, saline, complex, 0 to 2 percent slopes	5.70	4.15 %	30	5
F1167B 📃	Balaton-Wyard-Hamerly loams, 0 to 4 percent slopes	5.16	3.76 %	70	2
F1153B 📃	Souris-Balaton-Svea loams, 0 to 4 percent slope	4.82	3.51 %	79	2
F120A	Vallers, saline-Manfred complex, 0 to 1 percent slopes	3.88	2.83 %	33	4
F129B 📕	Cavour-Ferney loams, 3 to 6 percent slopes	3.71	2.70 %	30	4
F523A 📕	Lowe-Fluvaquents, channeled complex, 0 to 2 percent slopes, frequently flooded	2.78	2.03 %	19	6
			Average:	68.3	

Crop Productivity Index tables for ND and SD are obtained from SSURGO using weighted average of all components within the soil profile Soils data provided by USDA and NRCS.