






PLEGGENKUHLE (400.00 AC TOTAL, 302.63 AC TILLABLE)

# TOTAL SOILS



County: **Chickasaw County, IA**  
Location: **01-94N-11W**  
Township: **Fredericksburg**  
Acres: **400.00**  
Date: **05/22/2025**



Field borders obtained from Farm Service Agency  
as of 2008

FarmWorth, LLC makes no representations or warranties, express or implied, as to the accuracy of any information, data, numerical values, boundaries, or any other information generated through the use of FarmWorth.com. User is solely responsible for independently investigating and determining all information provided through FarmWorth.com prior to use and waives any and all claims against FarmWorth, LLC for any inaccuracies or inconsistencies in the information and/or data.

Code	Description	Acres	% of field	IA CSR2	Overall NCCPI	Soil Class (Irr)	Soil Class (Non-Irr)	Drainage Class
284B	Flagler sandy loam, 2 to 5 percent slopes	95.68	23.92 %	49	62	-	3	Somewhat excessively drained
84	Clyde clay loam, 0 to 3 percent slopes	75.69	18.92 %	88	90	-	2	Poorly drained
284C2	Flagler sandy loam, 5 to 9 percent slopes, moderately eroded	40.85	10.21 %	41	54	-	3	Somewhat excessively drained
198B	Floyd loam, 1 to 4 percent slopes	38.24	9.56 %	89	88	-	2	Somewhat poorly drained
391B	Clyde-Floyd complex, 1 to 4 percent slopes	30.66	7.66 %	87	89	-	2	Poorly drained
1152	Marshan clay loam, 0 to 2 percent slopes, rarely flooded	23.03	5.76 %	54	81	-	2	Poorly drained
394C2	Ostrander loam, 5 to 9 percent slopes, eroded	13.39	3.35 %	73	65	-	3	Well drained
621B	Houghton muck, 1 to 4 percent slopes	11.33	2.83 %	35	94	-	3	Very poorly drained
83B	Kenyon loam, 2 to 5 percent slopes	10.62	2.66 %	90	89	-	2	Moderately well drained
284C	Flagler sandy loam, 5 to 9 percent slopes	9.06	2.26 %	44	61	-	3	Somewhat excessively drained
407B	Schley loam, 1 to 4 percent slopes	8.52	2.13 %	81	94	-	2	Somewhat poorly drained
394C	Ostrander loam, 5 to 9 percent slopes	7.77	1.94 %	83	77	-	3	Well drained
221B	Klossner muck, 1 to 4 percent slopes	7.66	1.92 %	48	89	-	3	Very poorly drained
394B	Ostrander loam, 2 to 5 percent slopes	6.81	1.70 %	88	78	-	2	Well drained
171B	Bassett loam, 2 to 5 percent slopes	4.87	1.22 %	85	87	-	2	Moderately well drained
175B	Dickinson fine sandy loam, 2 to 5 percent slopes	4.46	1.12 %	50	61	-	3	Well drained
798B	Protivin loam, 1 to 4 percent slopes	3.81	0.95 %	61	79	-	2	Somewhat poorly drained
776D2	Lilah sandy loam, 5 to 14 percent slopes, moderately eroded	3.76	0.94 %	5	38	-	6	Excessively drained
775C	Billett sandy loam, 5 to 9 percent slopes	0.97	0.24 %	62	60	-	3	Well drained
399	Readlyn silt loam, 1 to 3 percent slopes	0.83	0.21 %	91	80	-	1	Somewhat poorly drained
83C2	Kenyon loam, 5 to 9 percent slopes, eroded	0.80	0.20 %	84	82	-	3	Moderately well drained
83C	Kenyon loam, 5 to 9 percent slopes	0.59	0.15 %	85	88	-	3	Moderately well drained
775B	Billett sandy loam, 2 to 5 percent slopes	0.50	0.12 %	67	61	-	3	Well drained
177	Saude loam, 0 to 2 percent slopes	0.10	0.02 %	60	78	-	2	Well drained
Average:				66.4	75.9			

IA has updated the CSR values for each county to CSR2.  
Soils data provided by USDA and NRCS.