





WEEHLER FARM (152.60 AC TOTAL, 105.00 AC TILLABLE)

TOTAL SOILS



County: Ringgold County, IA

Location: 01-67N-31W

Township: Clinton

Acres: 152.60















Date: 05/06/2025



Field borders obtained from Farm Service Agency
as of 2008

FarmWorth, LLC makes no representations or warranties, express or implied, as to the accuracy of any information, data, numerical values, boundaries, or any other information generated through the use of FarmWorth.com. User is solely responsible for independently investigating and determining all information provided through FarmWorth.com prior to use and waives any and all claims against FarmWorth, LLC for any inaccuracies or inconsistencies in the information and/or data.

Code	Description	Acres	% of field	IA CSR2	Overall NCCPI	Soil Class (Irr)	Soil Class (Non-Irr)	Drainage Class
S179F	Gara loam, 18 to 25 percent slopes	14.37	9.41 %	19	54	-	6	Well drained
179F2	Gara loam, 18 to 24 percent slopes, moderately eroded	14.19	9.30 %	12	54	-	6	Well drained
13B	Olmitz-Zook-Humeston complex, 0 to 5 percent slopes	14.15	9.27 %	78	77	-	2	Poorly drained
792D2	Armstrong clay loam, 9 to 14 percent slopes, moderately eroded	13.79	9.03 %	7	58	-	4	Somewhat poorly drained
179E2	Gara loam, 14 to 18 percent slopes, moderately eroded	12.33	8.08 %	24	62	-	6	Well drained
269	Humeston silty clay loam, 0 to 2 percent slopes, occasionally flooded	10.24	6.71 %	70	89	-	3	Poorly drained
175C	Dickinson fine sandy loam, 5 to 9 percent slopes	9.13	5.98 %	46	58	-	3	Well drained
76C	Ladoga silt loam, 5 to 9 percent slopes	8.13	5.33 %	81	88	-	3	Moderately well drained
S273C	Olmitz loam, heavy till, 5 to 9 percent slopes	7.89	5.17 %	77	84	-	3	Moderately well drained
451F2	Caleb loam, 14 to 25 percent slopes, moderately eroded	6.33	4.15 %	20	68	-	7	Moderately well drained
S993D2	Gara-Armstrong complex, 9 to 14 percent slopes, moderately eroded	6.27	4.11 %	32	64	-	4	Well drained
S220	Nodaway silt loam, heavy till, 0 to 2 percent slopes, occasionally flooded	5.92	3.88 %	77	86	-	2	Moderately well drained

Code		Description	Acres	% of field	IA CSR2	Overall NCCPI	Soil Class (Irr)	Soil Class (Non-Irr)	Drainage Class
S54		Zook silty clay loam, heavy till, 0 to 2 percent slopes, occasionally flooded	5.56	3.64 %	68	74	-	2	Poorly drained
179D		Gara loam, 9 to 14 percent slopes	4.15	2.72 %	42	70	-	4	Well drained
S54+		Zook silt loam, heavy till, 0 to 2 percent slopes, occasionally flooded, overwash	3.22	2.11 %	68	73	-	2	Poorly drained
175D		Dickinson fine sandy loam, 9 to 14 percent slopes	3.09	2.02 %	19	55	-	4	Well drained
94D2		Mystic-Caleb complex, 9 to 14 percent slopes, moderately eroded	2.92	1.91 %	20	72	-	4	Somewhat poorly drained
592D		Mystic loam, 9 to 14 percent slopes	2.60	1.70 %	10	67	-	4	Somewhat poorly drained
179E		Gara loam, 14 to 18 percent slopes	2.36	1.55 %	30	66	-	6	Well drained
792C		Armstrong loam, 5 to 9 percent slopes	2.24	1.47 %	35	66	-	3	Somewhat poorly drained
S172		Wabash silty clay, 0 to 2 percent slopes, occasionally flooded	1.34	0.88 %	57	62	-	3	Very poorly drained
W		Water	0.87	0.57 %	0	0	-	-	-
76C2		Ladoga silty clay loam, 5 to 9 percent slopes, moderately eroded	0.80	0.52 %	73	84	-	3	Moderately well drained
S269+		Humeston silt loam, 0 to 2 percent slopes, overwash, occasionally flooded	0.56	0.37 %	72	81	-	3	Poorly drained
269B		Humeston silty clay loam, 2 to 5 percent slopes, rarely flooded	0.17	0.11 %	71	89	-	3	Poorly drained
592C2		Mystic clay loam, 5 to 9 percent slopes, moderately eroded	0.03	0.02 %	31	70	-	3	Somewhat poorly drained
Average:					41.8	67.9			

IA has updated the CSR values for each county to CSR2.

Soils data provided by USDA and NRCS.